

15 November 2010
[23-10]

PROPOSAL M1006

MAXIMUM RESIDUE LIMITS (OCTOBER 2009- MARCH 2010)

ASSESSMENT REPORT

Executive Summary

Purpose

The purpose of this Proposal is to consider incorporating limits for residues of agricultural and veterinary chemicals that may legitimately occur in food in the *Australia New Zealand Food Standards Code* (the Code). This includes maximum residue limits (MRLs) gazetted by the Australian Pesticides and Veterinary Medicines Authority (APVMA) from October 2009 to March 2010. This Proposal also includes consideration of limits requested by other parties to further align the Code with international standards and other standards. This will permit the sale of foods containing legitimate residues and protect public health and safety by minimising residues in foods consistent with the effective control of pests and diseases.

Food Standards Australia New Zealand's (FSANZ's) role in the regulation of agricultural and veterinary chemicals is to protect public health and safety by ensuring that any potential residues in food are within appropriate safety limits, and to support industry and compliance agencies by maintaining limits in the Code that reflect legitimate residues in food.

Dietary exposure assessments indicate that in relation to current health-based guidance values, the proposed limits do not present any public health and safety concerns. This Proposal does not include consideration of any MRLs for antibiotic residues in food.

The *Agreement between the Government of Australia and the Government of New Zealand concerning a Joint Food Standards System* (the Treaty) excludes MRLs for agricultural and veterinary chemicals in food from the system setting joint food standards. Australia and New Zealand independently and separately develop MRLs for agricultural and veterinary chemicals in food.

FSANZ will make a Sanitary and Phytosanitary notification to the World Trade Organization (WTO).

Submissions are now invited on this Report to assist FSANZ finalise the assessment.

This Proposal is being assessed under the General Procedure.

Assessing the Proposal

In assessing the Proposal and the subsequent development of food regulatory measures, FSANZ has had regard to its statutory objectives in section 18 and the following matters prescribed in section 59 of the *Food Standards Australia New Zealand Act 1991* (FSANZ Act):

- Whether costs that would arise from a food regulatory measure developed or varied as a result of the Proposal outweigh the direct and indirect benefits to the community, Government or industry that would arise from the development or variation of the food regulatory measure
- There are no other measures that would be more cost-effective than a variation to Standard 1.4.2 that could achieve the same end
- Any relevant New Zealand standards
- Any other relevant matters.

Preferred Approach

To prepare draft variations to Standard 1.4.2 – Maximum Residue Limits.

Reasons for Preferred Approach

This Proposal has been assessed against the considerations provided for in section 59 of the FSANZ Act. FSANZ recommends the proposed draft variations to Standard 1.4.2 for the following reasons:

- MRLs serve to protect public health and safety by minimising residues in food consistent with the effective control of pests and diseases.
- Dietary exposure assessments indicate that the proposed variations do not present any public health and safety concerns.
- This approach ensures openness and transparency in relation to the residues that could reasonably occur in food.
- The proposed variations will benefit the community by maintaining public health and safety while permitting the legal sale of food containing legitimate residues of agricultural and veterinary chemicals used to control pests and diseases and improve agricultural productivity.
- The APVMA has assessed appropriate residue, animal transfer, processing and metabolism studies, in accordance with *The Manual of Requirements and Guidelines – MORAG – for Agricultural and Veterinary Chemicals 1 July 2005* to support the use of chemicals on commodities as outlined in this Proposal.
- The Office of Chemical Safety and Environmental Health (OCSEH) has undertaken a toxicological assessment of each chemical and has established an acceptable daily intake (ADI) and, where appropriate, an acute reference dose (ARfD).
- FSANZ has undertaken a preliminary regulation impact assessment and concluded that the proposed draft variations are necessary, cost-effective and beneficial.

- The proposed draft variations would remove inconsistencies between agricultural and food standards and provide certainty and consistency for producers, importers and Australian, State and Territory compliance agencies.
- The proposed changes are consistent with the FSANZ Act section 18 objectives.

Consultation

FSANZ is seeking public comment on this Assessment Report to assist in assessing the Proposal. Comments on, but not limited to, any impacts (costs/benefits) of the proposed variations, in particular the likely impacts on importation of food if the variations are advanced; any public health and safety considerations associated with the proposed limits; and any other affected parties would be welcome.

Invitation for Submissions

FSANZ invites public comment on this Report and the draft variations to the Code based on regulation impact principles for the purpose of preparing an amendment to the Code for approval by the FSANZ Board.

Written submissions are invited from interested individuals and organisations to assist FSANZ in further considering this Proposal. Submissions should, where possible, address the objectives of FSANZ as set out in section 18 of the FSANZ Act. Information providing details of potential costs and benefits of the proposed change to the Code from stakeholders is highly desirable. Claims made in submissions should be supported wherever possible by referencing or including relevant studies, research findings, trials, surveys etc. Technical information should be in sufficient detail to allow independent scientific assessment.

The processes of FSANZ are open to public scrutiny, and any submissions received will ordinarily be placed on the public register of FSANZ and made available for inspection. If you wish any information contained in a submission to remain confidential to FSANZ, you should clearly identify the sensitive information, separate it from your submission and provide justification for treating it as confidential commercial material. Section 114 of the FSANZ Act requires FSANZ to treat in-confidence, trade secrets relating to food and any other information relating to food, the commercial value of which would be, or could reasonably be expected to be, destroyed or diminished by disclosure.

Submissions must be made in writing and should clearly be marked with the word 'Submission' and quote the correct project number and name. While FSANZ accepts submissions in hard copy to our offices, it is more convenient and quicker to receive submissions electronically through the FSANZ website using the Changing the Code tab and then through Documents for Public Comment. Alternatively, you may email your submission directly to the Standards Management Officer at submissions@foodstandards.gov.au. There is no need to send a hard copy of your submission if you have submitted it by email or the FSANZ website. FSANZ endeavours to formally acknowledge receipt of submissions within 3 business days.

DEADLINE FOR PUBLIC SUBMISSIONS: 6pm (Canberra time) 13 December 2010

SUBMISSIONS RECEIVED AFTER THIS DEADLINE WILL NOT BE CONSIDERED

Submissions received after this date will only be considered if agreement for an extension has been given prior to this closing date. Agreement to an extension of time will only be given if extraordinary circumstances warrant an extension to the submission period. Any agreed extension will be notified on the FSANZ website and will apply to all submitters.

Questions relating to making submissions or the application process can be directed to the Standards Management Officer at standards.management@foodstandards.gov.au.

If you are unable to submit your submission electronically, hard copy submissions may be sent to one of the following addresses:

Food Standards Australia New Zealand
PO Box 7186
Canberra BC ACT 2610
AUSTRALIA
Tel (02) 6271 2222

Food Standards Australia New Zealand
PO Box 10559
The Terrace WELLINGTON 6036
NEW ZEALAND
Tel (04) 978 5636

CONTENTS

INTRODUCTION	2
1. THE ISSUE / PROBLEM.....	3
2. CURRENT STANDARD.....	3
2.1 <i>Background</i>	3
3. OBJECTIVES	3
4. ASSESSMENT APPROACH.....	4
RISK ASSESSMENT.....	4
5. RISK ASSESSMENT SUMMARY	4
RISK MANAGEMENT	5
6. OPTIONS	5
7. IMPACT ANALYSIS	5
7.1 <i>Affected Parties</i>	5
7.2 <i>Benefit Cost Analysis</i>	5
7.3 <i>Comparison of Options</i>	7
COMMUNICATION AND CONSULTATION STRATEGY.....	8
8. COMMUNICATION	8
9. CONSULTATION.....	8
9.1 <i>World Trade Organization (WTO)</i>	8
9.2 <i>Codex Alimentarius Commission Standards</i>	9
9.3 <i>New Zealand Standards</i>	11
9.4 <i>Imported foods</i>	13
CONCLUSION.....	14
10. CONCLUSION AND PREFERRED OPTION	14
10.1 <i>Reasons for Preferred Approach</i>	14
11. IMPLEMENTATION AND REVIEW	14
ATTACHMENT 1 – DRAFT VARIATIONS TO THE <i>AUSTRALIA NEW ZEALAND FOOD STANDARDS</i> <i>CODE</i>	16
ATTACHMENT 2 – SUMMARY OF PROPOSED MRLS AND TECHNICAL AMENDMENTS IN PROPOSAL M1006.....	23

SUPPORTING DOCUMENTS

The following documents are available on the FSANZ website at:

http://www.foodstandards.gov.au/foodstandards/proposals/proposal_m1006_maximum4786.cfm

- SD1: Safety Assessment Methodology
- SD2: Background Information.

Introduction

Notifications were received from the Australian Pesticides and Veterinary Medicines Authority (APVMA) on 2 October, 4 November and 8 December 2009 and 5 and 8 February, 10 March and 12 April 2010 seeking to vary the *Australia New Zealand Food Standards Code* (the Code). The proposed variations to the Code would align maximum residue limits (MRLs) in the Code for certain agricultural and veterinary chemicals with the APVMA MRLs listed in *The MRL Standard* and permit the sale of relevant foods containing legitimate residues.

This Proposal also includes consideration of varying MRLs for bifenazate, buprofezin, carbaryl, chlorpyrifos, cypermethrin, fenbuconazole, fenbutatin oxide, lambda-cyhalothrin, metconazole (new entry), methoxyfenozide, profenofos, spirotetramat, tebuconazole, tebufenozide and trifloxystrobin as a result of information provided by other parties. Anomalies between the Code and international standards may have implications for trade in certain foods. The proposed variations would align limits in the Code with Codex and other countries' standards and permit the sale of relevant foods containing legitimate residues at levels that do not present health or safety concerns.

In summary, this Proposal includes consideration of MRL variations for abamectin, benzyladenine, beta-cyfluthrin, bifenazate, bifenthrin, boscalid, bromoxynil, buprofezin, carbaryl, chlorothalonil, chlorpyrifos, clothianidin, cypermethrin, epoxiconazole, etoxazole, fenbuconazole, fenbutatin oxide, fipronil, fluazifop-butyl, flubendiamide, flumetsulam, imazamox, imazapyr, imidacloprid, indoxacarb, iprodione, lambda-cyhalothrin, metalaxyl, metalaxyl-M, metconazole, methomyl, methoxyfenozide, paclobutrazol, pendimethalin, permethrin, phosphorous acid, pirimicarb, profenofos, prothioconazole, pyraclostrobin, pyrimethanil, pyriproxyfen, simazine, spirotetramat, tebuconazole, tebufenozide, terbutylazine, tolclofos-methyl, triadimenol, trichlorfon, trifloxystrobin, trifluralin and trinexapac-ethyl.

The draft variations to the Code are at **Attachment 1**. An outline of these variations and dietary exposure estimates is at **Attachment 2**. The safety assessment methodology is outlined in **Supporting Document 1**. This includes an explanation of terminology.

FSANZ's role in the regulation of agricultural and veterinary chemicals is to protect public health and safety by ensuring that any potential residues in food are within appropriate safety limits, and to support producers, importers and compliance agencies by maintaining limits in the Code that reflect legitimate residues in food.

In considering the issues associated with variations to limits in the Code for residues of agricultural and veterinary chemicals in food, it should be noted that the limit is the maximum level of a chemical that may be in a food, not the level that is usually present in a food. However, incorporating the limit into food legislation means that the residues of a chemical are minimised (i.e. must not exceed the MRL or other limit), irrespective of whether the dietary exposure assessment indicates that higher residues would not risk public health and safety.

Limits and variations to limits in the Code do not permit or prohibit the use of agricultural or veterinary chemicals. Other Australian Government, State and Territory legislation regulates use and control of agricultural and veterinary chemicals.

1. The Issue / Problem

Including limits for residues of agricultural and veterinary chemicals in foods in the Code has the effect of allowing the sale of food containing legitimate residues, where any residues do not exceed these limits. Variations in MRLs reflect the changing use patterns of agricultural and veterinary chemicals available to chemical product users including food producers. These changes include both the development of new products and crop uses, and the withdrawal of older products following review. Where residues do not pose health or safety concerns, limits are also varied in line with international standards to reflect requirements for foods containing legitimate residues to be imported. Internationally, farmers face different pest and disease pressures and so agricultural and veterinary chemical use patterns may vary.

2. Current Standard

2.1 Background

Standard 1.4.2 lists the limits for agricultural and veterinary chemical residues which may occur in foods. If a limit is not listed for a particular agricultural or veterinary chemical/food combination, there must be no detectable residues of that chemical in that food. This general prohibition means that in the absence of the relevant limit in the Code, food may not be sold where there are detectable residues.

Variations to the Code may be required to permit the sale of foods containing legitimate residues. A dietary exposure assessment is conducted before the Code is varied to ensure that proposed limits do not present any public health or safety concerns.

Further background information on MRLs, the regulatory framework for agricultural and veterinary chemicals and the FSANZ assessment process for incorporating limits, including MRLs for antibiotic substances, in the Code is provided in **Supporting Document 2**.

3. Objectives

In assessing this Proposal, FSANZ aims to ensure that approving the proposed draft variations does not present public health and safety concerns and that the sale of food containing legitimate residues is permitted.

In developing or varying a food standard, FSANZ is required by its legislation to meet three primary objectives which are set out in section 18 of the FSANZ Act. These are:

- the protection of public health and safety; and
- the provision of adequate information relating to food to enable consumers to make informed choices; and
- the prevention of misleading or deceptive conduct.

In developing and varying standards, FSANZ must also have regard to:

- the need for standards to be based on risk analysis using the best available scientific evidence;
- the promotion of consistency between domestic and international food standards;

- the desirability of an efficient and internationally competitive food industry;
- the promotion of fair trading in food; and
- any written policy guidelines formulated by the Ministerial Council.

4. Assessment Approach

FSANZ's primary role in developing food regulatory measures for residues of agricultural and veterinary chemicals in food is to ensure that the potential residues are within health-based guidance values. FSANZ conducts and reviews dietary exposure assessments in accordance with internationally accepted practices and procedures.

In assessing the public health and safety implications of chemical residues, FSANZ considers the dietary exposure to chemical residues from potentially treated foods in the diet by comparing the dietary exposure with the relevant health-based guidance value. FSANZ will not approve variations to limits in the Code where dietary exposure to the residues of a chemical could risk public health and safety.

The steps undertaken in conducting a dietary exposure assessment are:

- determining the residues of a chemical in a treated food; and
- calculating the dietary exposure to a chemical from relevant foods, using food consumption data from national nutrition surveys and comparing this to the relevant health-based guidance value.

The estimated dietary exposure to a chemical is compared to the relevant health-based guidance value/s for that chemical in food (i.e. the acceptable daily intake (ADI) and/or the acute reference dose (ARfD)). FSANZ considers that dietary exposure to the residues of a chemical is acceptable where the best estimate of this exposure does not exceed the relevant guidance value/s.

The safety assessment methodology is further outlined in **Supporting Document 1**.

RISK ASSESSMENT

5. Risk Assessment Summary

FSANZ has reviewed the dietary exposure assessments submitted by the APVMA and conducted dietary exposure assessments to assess the limits requested by other parties. Using the best available scientific data and internationally recognised risk assessment methodology, FSANZ concluded that in relation to current health-based guidance values, setting the limits as proposed does not present any public health and safety concerns.

The additional safety factors inherent in calculation of the ADI and ARfD mean that there is negligible risk to public health and safety when estimated exposures are below these guidance values.

Risk Management

6. Options

After the submission period, the following options are available:

1. Option 1 – approve the draft variations
2. Option 2 – approve the draft variations subject to such amendments as FSANZ considers necessary
3. Option 3 – abandon the proposal

7. Impact Analysis

The impact analysis represents likely impacts based on available information. The impact analysis is designed to assist in the process of identifying affected parties and any alternative options consistent with the objective of the proposed changes. Information from public submissions is sought to further assess the proposed changes.

The draft variations may be amended and option 2 recommended for approval where the need is identified. For example, an MRL may be retained rather than reduced or deleted where the necessity for the MRL to allow for the importation and sale of safe food is identified through consultation. Further information to assist in identifying implications for imported foods is provided in section 9 of this Report and the requested variations are presented in **Attachments 1 and 2**.

7.1 Affected Parties

The sectors of the community potentially affected by the proposed amendments include:

- consumers
- growers and producers
- importers of agricultural produce and food products
- the chemical industry
- Australian and New Zealand Government, State and Territory agencies involved in monitoring and regulating the use of agricultural and veterinary chemicals in food and the potential resulting residues.

7.2 Benefit Cost Analysis

7.2.1 Option 1 – approve the draft variations

This option may contribute to community confidence that regulatory authorities are maintaining standards to minimise residues of agricultural and veterinary chemicals in the food supply. FSANZ does not consider there to be any adverse dietary exposure implications associated with the proposed approval. The risk assessment has determined that there are no public health or safety concerns associated with the proposed variations. No additional costs to consumers have been identified.

This option benefits growers and producers in Australia as agricultural and food standards are further aligned. This means that foods produced in accordance with agricultural standards and legislation may be sold under food legislation as MRL variations are incorporated in the Code. The proposed variations are unlikely to result in any costs for producers as changes in use patterns are made as required; current proper use results in compliance with the proposed variations already.

Importers may benefit or be disadvantaged by the approval of the proposed draft variations. Additional or increased MRLs may benefit importers and consequently consumers in that this may extend the options to source safe foods. Any MRL deletions or reductions have the potential to restrict importation of foods and could potentially result in higher food prices and a reduced product range available to consumers.

This option benefits Australian Government, State and Territory agencies in that it serves to further harmonise agricultural and food standards. This is of particular assistance to compliance agencies. Achieving further consistency between agricultural and food legislation would minimise compliance costs to primary producers and assist in efficient enforcement of regulations. This option is unlikely to result in discernable costs to Government agencies, although an awareness of changes in the standards for residues in food would be needed and there may be minimal impacts associated with slight changes to residue monitoring programs.

Interested parties are invited to comment on any impacts of the proposed variations during the public consultation period. This is to ensure that any adverse consequences of the proposed variations can be addressed. Imported foods and Codex MRLs are addressed in section 9 of this Report.

7.2.2 Option 2 – approve the draft variations subject to such amendments as FSANZ considers necessary

FSANZ will consider any comments received during the submission period and may amend the draft variations following further assessment.

7.2.3 Option 3 – abandon the Proposal

This option would allow inconsistencies between agricultural and food legislation to perpetuate as the Code would not reflect residues that may be present in foods following legitimate use of chemical products in Australia as determined by the APVMA. This may result in foods legitimately treated during production not being permitted for sale. Producers would incur significant costs. This may also create uncertainty, inefficiency and confusion in the enforcement of regulations. In addition, the anomalies between the Code and international standards identified by industry would perpetuate and may have implications for trade in certain foods. This would impact negatively on all affected parties and producers, industry and compliance agencies in particular.

Importers may benefit if proposed MRL deletions or reductions are not progressed as the continuity of existing limits could be relied upon. However, there is scope under current processes to retain specific MRLs where the necessity for the MRL to continue to allow the importation and sale of safe food is identified through consultation. This is discussed in section 9 of this Report. Importers and consequently consumers may be disadvantaged where proposed additional or increased MRLs are not progressed as this may unnecessarily limit sources of certain foods.

7.2.4 Summary

FSANZ conducted a Best Practice Regulation Preliminary Assessment and concluded that business compliance costs and other impacts on business, individuals, regulatory agencies and the economy are low or nil. The regulatory proposal does not impose impacts on business, individuals, regulatory agencies or the economy that warrant further analysis. The changes to regulation are machinery in nature involving technical variations to the Standard which will not have appreciable impacts and are consistent with existing policy.

FSANZ consulted with the Office of Best Practice Regulation (OBPR) on the need for the preparation of a regulation impact statement (RIS) under the Council of Australian Governments' requirements. The OBPR concluded that the proposed changes are minor and do not substantially alter existing arrangements. The OBPR advised that a RIS is therefore not required.

7.3 Comparison of Options

In assessing proposed variations to the Code, FSANZ considers the impact of various regulatory and non-regulatory options on all sectors of the community, including consumers, food industries and governments in Australia.

FSANZ recommends approving option 1 – approve the draft variations for the following reasons:

- There are no public health and safety concerns associated with the proposed variations.
- This approach ensures openness and transparency in relation to the residues that could reasonably occur in food.
- The changes would minimise potential costs to primary producers, rural and regional communities and importers in terms of permitting the sale of food containing legitimate residues.
- The changes would minimise residues in food consistent with the effective use of agricultural and veterinary chemicals to control pests and diseases.
- The changes would further align the Code with international standards.
- The changes would remove inconsistencies between agricultural and food standards and assist compliance agencies.

Option 2 may be recommended at the Approval stage subject to the need for any required amendments being identified through consultation and further assessment.

Option 3 is an undesirable option because potential substantial costs to primary producers may result. Additional costs may impact negatively on their viability and in turn the viability of the rural and regional communities that depend upon the sale of agricultural produce. This option may restrict the opportunity for importers to source safe produce or foods internationally and potentially impact consumers through higher food prices and limited choice. Also, consequent inconsistencies between agricultural and food legislation could have negative impacts on compliance costs for producers, perception problems in export markets and undermine the efficient enforcement of standards for chemical residues.

The benefits of progressing option 1 outweigh any associated costs.

Communication and Consultation Strategy

8. Communication

Consideration of amending limits in the Code for residues of agricultural or veterinary chemicals in food does not normally generate public interest. FSANZ adopts a basic communication strategy, with a focus on alerting the community that changes to the Code are being contemplated.

FSANZ publishes the details of proposed changes and subsequent reports on its website (<http://www.foodstandards.gov.au/foodstandards/proposals/proposalm1006maximum4786.cfm>), alerts subscribers (over 5000) via email of the availability of these reports for comment, and issues media releases drawing attention to proposed Code amendments.

The Applicant, individuals and organisations making submissions on this Application will be notified at each stage of the Application. If the FSANZ Board approves the draft variation to the Code, FSANZ will notify its decision to the Ministerial Council. The Applicant and stakeholders, including the public, will be notified of the gazetted changes to the Code in the national press and on the FSANZ website.

Once the Code has been amended, FSANZ incorporates the changes in the website version of the Code and, through its email and telephone information service, responds to community enquiries.

Should the media show an interest in any of the assessed chemicals, FSANZ or the APVMA can provide background information as required.

9. Consultation

FSANZ is seeking public comment on the proposed changes to the Code outlined in this Report to assist in finalising the assessment. Comments on, but not limited to, any impacts (costs/benefits) of the proposed variations, in particular the likely impacts on importation of food if specific variations are advanced; any public health and safety considerations associated with the proposed changes; and any other affected parties would be useful.

9.1 World Trade Organization (WTO)

As a member of the World Trade Organization (WTO), Australia is obligated to notify WTO member nations where proposed mandatory regulatory measures are inconsistent with any existing or imminent international standards and the proposed measure may have a significant effect on trade.

Limits prescribed in the Code constitute a mandatory requirement applying to all food products of a particular class whether produced domestically or imported. Food products with residues exceeding the relevant limit listed in the Code cannot legally be supplied in Australia.

This Proposal includes consideration of varying limits in the Code for residues of agricultural and veterinary chemicals in food that are addressed in the international Codex standard. Limits in the Proposal relate to chemical residues that may occur in heavily traded agricultural commodities that may indirectly have a significant effect on trade of derivative food products between WTO members.

This Proposal will be notified as a Sanitary and Phytosanitary (SPS) measure in accordance with the WTO Agreement on the Application of SPS Measures as the primary objective of the measure is to support the regulation of the use of agricultural and veterinary chemical products to protect human, animal and plant health and the environment.

9.2 Codex Alimentarius Commission Standards

Codex standards are used as the relevant international standard to determine whether a new or changed standard requires a WTO notification.

FSANZ may consider varying limits for residues of agricultural or veterinary chemicals in food in a Proposal where interested parties have identified anomalies between the Code and international standards that may result in adverse impacts. FSANZ must have regard to its WTO obligations, the promotion of consistency between domestic and international food standards and the promotion of fair trading in food. These matters encompass a consideration of international standards and trade issues. The assessment gives careful consideration to public health and safety.

Interested parties provided information that specific anomalies between the Code and Codex or other standards may present barriers to trade in certain foods. This Proposal includes proposed limits for bifenazate, buprofezin, carbaryl, chlorpyrifos, cypermethrin, fenbuconazole, fenbutatin oxide, lambda-cyhalothrin, metconazole, methoxyfenozide, profenofos, spirotetramat, tebuconazole, tebufenozide and trifloxystrobin to address these issues. Further detail is provided at **Attachment 2**. The proposed variations to the Code would align limits in the Code with international standards or standards in producer or other importing countries and permit the sale of relevant foods containing legitimate residues that do not present health or safety concerns.

The following table lists proposed limits where there is a corresponding Codex limit.

Chemical Food	Proposed limit ^{†‡} mg/kg	Codex limit mg/kg
Bifenazate Stone fruits [except plums]	2.5	Stone fruits 2
Boscalid Edible offal (mammalian) Brassica leafy vegetables Lettuce, head Lettuce leaf Meat (mammalian) (in the fat) Milks	0.3 T30 T15 T15 0.3 0.1	0.2 Leafy vegetables 30 Meat (from mammals other than marine mammals) (fat) 0.7 0.1
Carbaryl Cranberry	3	5
Chlorothalonil Herbs Pulses	T20 3	Celery leaves 3 Parsley 3 Beans (dry) 0.2
Chlorpyrifos Cranberry	1	1
Cyhalothrin Stone fruits	0.5	Apricot 0.5 Cherries 0.3 Nectarine 0.5 Peach 0.5 0.2 Plums (including prunes)

Chemical Food	Proposed limit ^{†‡} mg/kg	Codex limit mg/kg
Cypermethrin Durian Longan Peppers, Chili	1 1 1	*1 1 Peppers, Chili, dried 2
Fenbuconazole Edible offal (mammalian) Stone fruits [except nectarine] Wheat	0.05 1 *0.01	0.1 Apricot 0.5 Cherries 1 Peach 0.5 0.1
Fenbutatin oxide Cherries	6	10
Fipronil Sweet potato	*0.01	Potato 0.02
Imidacloprid Field pea (dry) Leafy vegetables [except lettuce, head] Lettuce, head Potato Sweet potato	T*0.05 20 5 0.3 0.3	Peas (dry) 2 Radish leaves (including radish tops) 5 2 Root and tuber vegetables 0.5
Indoxacarb Peanut	T0.02	*0.02
Methoxyfenozide Cranberry Stone fruits [except plums]	0.5 3	0.7 Stone fruits 2
Pirimicarb Adzuki bean (dry) Mung bean (dry) Leafy vegetables [except chervil; mizuna; rucola (rocket)]	T0.5 T0.5 T7	Pulses 0.2 Kale 0.3 Lettuce, Head 5 Lettuce, Leaf 5
Profenofos Mangosteen	5	10
Prothioconazole Barley Edible offal (mammalian) Oats Wheat	0.3 0.1 *0.05 0.3	0.2 0.5 0.05 0.1
Pyraclostrobin Cereal grains Papaya (pawpaw)	*0.01 T0.5	Barley 0.5 Maize *0.02 Oats 0.5 Spelt 0.2 Wheat 0.2 Papaya *0.05
Pyrimethanil Leafy vegetables	T5	Lettuce, Head 3

Chemical Food	Proposed limit^{†‡} mg/kg	Codex limit mg/kg
Spirotetramat		
Citrus fruits	1	0.5
Dried grapes	4	Dried grapes (=currants, raisins and sultanas) 4
Fruiting vegetables, other than cucurbits	7	1
Fruiting vegetables, cucurbits [except melons]	2	Fruiting vegetables, Cucurbits 0.2
Grapes	2	2
Melons, except watermelon	0.5	Fruiting vegetables, Cucurbits 0.2
Watermelon	0.5	Fruiting vegetables, Cucurbits 0.2
Leafy vegetables [except lettuce, head]	5	Leafy vegetables 7
Lettuce, head	3	
Potato	5	0.8
Tebuconazole		
Cherries	5	5
Tebufenozide		
Cranberry	0.5	0.5
Triadimenol		
Peppers	T1	Fruiting vegetables other than cucurbits 1
Peppers, Sweet	T1	Peppers, Sweet (including pimento or pimienta) 0.1
Trifloxystrobin		
Celery	T1	1
Stone fruits	2	3

[†] Note that a 'T' indicates that the limit is temporary.

[‡] An asterisk indicates that the limit is at or about the limit of analytical quantification.

FSANZ invites comment on any possible ramifications of approving the proposed MRLs.

9.3 New Zealand Standards

All imported and domestically produced food sold in New Zealand (except for food imported from Australia) must comply with the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards 2010 and amendments (the New Zealand MRL Standards).

Under the New Zealand MRL Standards, agricultural chemical residues in food must comply with the specific MRLs listed in the Standards. The New Zealand MRL Standards also include a provision for residues of up to 0.1 mg/kg for agricultural chemical / commodity combinations not specifically listed. If the food is imported, it may comply with Codex MRLs. Further information about the New Zealand MRL Standards is available on the New Zealand Food Safety Authority website at <http://www.nzfsa.govt.nz/registers-lists/nz-mrl/>.

Limits in the Code and in the New Zealand MRL Standards may differ for a number of legitimate reasons including differing use patterns for chemical products as a result of varying pest and disease pressures and varying climatic conditions.

The following table lists proposed MRLs where there is a corresponding MRL in the New Zealand Standards.

Chemical Food	Proposed MRL[†] mg/kg	NZ MRL[‡] mg/kg
Bifenthrin Fruiting vegetables, cucurbits [except cucumber]	0.1	Pumpkins *0.001 Squash *0.001
Carbaryl Cranberry	3	Fruits 3
Chlorothalonil Leafy vegetables [except chard (silver beet); spinach]	T10	Lettuce 10
Chlorpyrifos Blueberries Cherries Cranberry Stone fruits [except cherries]	*0.01 1 1 T1	Fruits (except bananas, grapes, kiwifruit and stone fruits) 0.2 Stone fruits 1
Epoxiconazole Cereal grains	0.05	Barley *0.05 Wheat *0.05
Fenbutatin oxide Cherries	6	Stone fruits 1
Imidacloprid Lettuce, head Potato Sweet potato	5 0.3 0.3	Lettuce 1 Potatoes *0.02
Pirimicarb Adzuki bean (dry) Mung bean(dry) Leafy vegetables [except chervil; mizuna; rucola (rocket)]	T0.5 T0.5 T7	Legume vegetables 0.5 Leafy vegetables 1
Prothioconazole Barley Cereal bran, unprocessed Oats Wheat Wheat germ	0.3 0.5 *0.05 0.3 0.5	Cereal grains *0.02
Pyraclostrobin Cereal grains	*0.01	Barley *0.02 Wheat *0.02
Spirotetramat Potato Sweet potato	5 5	Potatoes 0.5
Tebuconazole Cherries	5	Stone fruits 1
Trifloxystrobin Stone fruits	2	Stone fruits (except cherries) *0.02
Trinexapac-ethyl Barley Wheat	T0.3 T0.3	Cereal grains *0.05

[†] Note that a 'T' indicates that the limit is temporary.

[‡] An asterisk indicates that the limit is at or about the limit of analytical quantification.

FSANZ requests comment on the proposed MRLs in relation to the corresponding New Zealand MRLs.

9.4 Imported foods

Internationally, countries set MRLs according to good agricultural practice (GAP) or good veterinary practice (GVP). Agricultural and veterinary chemicals are used differently in different countries around the world as pests, diseases and environmental factors differ and because product use patterns may differ. This means that residues in imported foods may be legitimately different from those in domestically produced foods.

Deletions or reductions of MRLs may impact imported foods that may comply with existing MRLs even though these existing MRLs are no longer required for domestically produced food. This is because imported foods may contain residues consistent with the MRLs proposed for deletion or reduction.

FSANZ is committed to ensuring that the implications of MRL variations are considered. Under the current process for considering variations to the Code, FSANZ encourages submissions including specific data demonstrating a need for certain MRLs to be varied. FSANZ will consider amending proposed MRL variations to continue to allow the sale of safe food where such MRLs are supported by adequate data or information demonstrating that the residues are legitimate and likely to occur. The assessment will consider dietary exposure in the context of the Australian diet. Further information on data requirements may be obtained from FSANZ.

To assist in identifying possible impacts on imported foods, FSANZ has compiled the following table of foods where the MRLs are proposed for deletion or reduction. The proposed MRL variations to the Code are at **Attachment 1** and the proposed changes are outlined in **Attachment 2**.

Chemical Food
Chlorothalonil Pulses
Chlorpyrifos Blueberries
Imidacloprid Potato
Iprodione Brussels sprouts
Metalaxyl Papaya (pawpaw)
Pirimicarb Adzuki bean (dry) Mung bean (dry)
Spirotetramat Lettuce, head Lettuce, leaf Melons, except watermelon Watermelon
Tolclofos-methyl Beetroot

FSANZ requests comment on any possible ramifications for imported foods of the proposed variations.

Conclusion

10. Conclusion and Preferred Option

This Proposal has been assessed against the considerations provided for in section 59 of the FSANZ Act.

Preferred Approach

To prepare draft variations to Standard 1.4.2 – Maximum Residue Limits.

10.1 Reasons for Preferred Approach

FSANZ recommends the proposed draft variations to Standard 1.4.2 for the following reasons:

- MRLs serve to protect public health and safety by minimising residues in food consistent with the effective control of pests and diseases.
- Dietary exposure assessments indicate that the proposed variations do not present any public health and safety concerns.
- This approach ensures openness and transparency in relation to the residues that could reasonably occur in food.
- The proposed variations will benefit the community by maintaining public health and safety while permitting the legal sale of food with legitimate residues of agricultural and veterinary chemicals used to control pests and diseases and improve agricultural productivity.
- The APVMA has assessed appropriate residue, animal transfer, processing and metabolism studies, in accordance with *The Manual of Requirements and Guidelines – MORAG – for Agricultural and Veterinary Chemicals 1 July 2005* to support the use of chemicals on commodities as outlined in this Proposal.
- The OCSEH has undertaken a toxicological assessment of each chemical and has established an ADI and, where appropriate, an ARfD.
- FSANZ has undertaken a preliminary regulation impact assessment and concluded that the proposed draft variations are necessary, cost-effective and beneficial.
- The proposed draft variations would remove inconsistencies between agricultural and food standards and provide certainty and consistency for producers, importers and Australian, State and Territory compliance agencies.
- The proposed changes are consistent with the FSANZ Act section 18 objectives.

11. Implementation and Review

The use of chemical products and MRLs are under constant review as part of the APVMA Chemical Review Program. In addition, regulatory agencies continue to monitor health, agricultural and environmental issues associated with chemical product use.

Residues in food are also monitored through:

- State and Territory residue monitoring programs
- Australian Government programs such as the National Residue Survey
- dietary exposure studies such as the Australian Total Diet Study.

These monitoring programs and the continual review of the use of agricultural and veterinary chemicals mean that there is considerable scope to review limits in the Code.

It is proposed that the variations in this Proposal should take effect on gazettal and that the limits be subject to existing monitoring arrangements.

ATTACHMENTS

1. Draft variations to the *Australia New Zealand Food Standards Code*
2. Summary of proposed MRLs and technical amendments in Proposal M1006

Attachment 1

Draft variations to the *Australia New Zealand Food Standards Code*

Subsection 94 of the FSANZ Act provides that standards or variations to standards are legislative instruments, but are not subject to disallowance or sunseting

To commence: on gazettal

[1] **Standard 1.4.2** of the *Australia New Zealand Food Standards Code* is varied by –

[1.1] *omitting from Schedule 1 the chemical residue definition for the chemical appearing in Column 1 of the Table to this sub-item, substituting the chemical residue definition appearing in Column 2 –*

COLUMN 1	COLUMN 2
CHLOROTHALONIL	COMMODITIES OF PLANT ORIGIN: CHLOROTHALONIL COMMODITIES OF ANIMAL ORIGIN: 4-HYDROXY- 2,5,6-TRICHLOROISOPHTHALONITRILE METABOLITE, EXPRESSED AS CHLOROTHALONIL

[1.2] *inserting in Schedule 1 –*

METCONAZOLE METCONAZOLE	
STONE FRUITS	0.2

[1.3] *omitting from Schedule 1 the foods and associated MRLs for each of the following chemicals –*

BIFENTHRIN BIFENTHRIN	
FRUITING VEGETABLES, CUCURBITS	0.1
CHLOROTHALONIL COMMODITIES OF PLANT ORIGIN: CHLOROTHALONIL COMMODITIES OF ANIMAL ORIGIN: SUM OF CHLOROTHALONIL AND 4-HYDROXY-2, 5, 6- TRICHLOROISOPHTHALONITRILE METABOLITE, EXPRESSED AS CHLOROTHALONIL	
LEAFY VEGETABLES	T7
VEGETABLES [EXCEPT AS OTHERWISE LISTED UNDER THIS CHEMICAL]	T7
CHLORPYRIFOS CHLORPYRIFOS	
STONE FRUITS	T1
EPOXICONAZOLE EPOXICONAZOLE	
BARLEY	0.05

WHEAT	0.05
IMIDACLOPRID	
SUM OF IMIDACLOPRID AND METABOLITES CONTAINING THE 6-CHLOROPYRIDINYLMETHYLENE MOIETY, EXPRESSED AS IMIDACLOPRID	
LEAFY VEGETABLES [EXCEPT LETTUCE, LEAF]	T5
LETTUCE, LEAF	T20
PERMETHRIN	
PERMETHRIN, SUM OF ISOMERS	
CORIANDER (LEAVES AND STEMS)	T10
PIRIMICARB	
SUM OF PIRIMICARB, DEMETHYL-PIRIMICARB AND THE <i>N</i> -FORMYL-(METHYLAMINO) ANALOGUE (DEMETHYLFORMAMIDO-PIRIMICARB), EXPRESSED AS PIRIMICARB	
LEAFY VEGETABLES [EXCEPT CHERVIL; MIZUNA; RUCOLA]	T5
VEGETABLES [EXCEPT LEAFY VEGETABLES; LUPIN (DRY); SOYA BEAN (DRY); SWEET CORN (CORN-ON-THE-COB)]	1
SPIROTETRAMAT	
SUM OF SPIROTETRAMAT, AND CIS-3-(2,5- DIMETHYLPHENYL)-4-HYDROXY-8-METHOXY-1- AZASPIRO[4.5]DEC-3-EN-2-ONE, EXPRESSED AS SPIROTETRAMAT	
FRUITING VEGETABLES, CUCURBITS	T2
LETTUCE, LEAF	T10
PEPPERS, SWEET	T5
TOMATO	T7
TRIADIMENOL	
TRIADIMENOL SEE ALSO TRIADIMEFON	
PEPPERS, SWEET	T1
TRICHLORFON	
TRICHLORFON	
FRUIT [EXCEPT AS OTHERWISE LISTED UNDER THIS CHEMICAL]	0.1
VEGETABLES [EXCEPT AS OTHERWISE LISTED UNDER THIS CHEMICAL]	0.1

[1.4] inserting in alphabetical order in Schedule 1, the foods and associated MRLs for each of the following chemicals –

ABAMECTIN	
SUM OF AVERMECTIN B1A, AVERMECTIN B1B AND (Z)-8,9 AVERMECTIN B1A, AND (Z)-8,9 AVERMECTIN B1B	
SWEET CORN (CORN-ON-THE-COB)	T*0.01
BENZYLADENINE	
BENZYLADENINE	
PISTACHIO NUT	T*0.05
BIFENAZATE	
SUM OF BIFENAZATE AND BIFENAZATE DIAZENE (DIAZENECARBOXYLIC ACID, 2-(4-METHOXY-[1,1'-BIPHENYL-3-YL] 1-METHYLETHYL ESTER), EXPRESSED AS BIFENAZATE	
CHERRIES	2.5
LETTUCE, HEAD	T5
LETTUCE, LEAF	T5
BIFENTHRIN	
BIFENTHRIN	
CUCUMBER	T0.3
FRUITING VEGETABLES, CUCURBITS [EXCEPT CUCUMBER]	0.1
PINEAPPLE	T*0.01
BOSCALID	
COMMODITIES OF PLANT ORIGIN: BOSCALID COMMODITIES OF ANIMAL ORIGIN: SUM OF BOSCALID, 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE AND THE GLUCURONIDE CONJUGATE OF 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE, EXPRESSED AS BOSCALID EQUIVALENTS	
ALL OTHER FOODS	0.5
MILK FATS	0.7
BUPROFEZIN	
BUPROFEZIN	
STONE FRUITS [EXCEPT APRICOT; PEACH]	1.9
CARBARYL	
CARBARYL	
CRANBERRY	3
CHLOROTHALONIL	
COMMODITIES OF PLANT ORIGIN: CHLOROTHALONIL COMMODITIES OF ANIMAL ORIGIN: SUM OF CHLOROTHALONIL AND 4-HYDROXY-2, 5, 6-TRICHLOROISOPHTHALONITRILE METABOLITE, EXPRESSED AS CHLOROTHALONIL	
CHARD (SILVER BEET)	T50
CORIANDER (LEAVES, STEM, ROOTS)	T20

LEAFY VEGETABLES [EXCEPT CHARD (SILVER BEET); SPINACH]	T10
POULTRY, EDIBLE OFFAL OF	*0.05
POULTRY MEAT	*0.05
SPINACH	T100
VEGETABLES [EXCEPT ASPARAGUS; BRUSSELS SPROUTS; CARROT; CELERY; CHARD (SILVER BEET); FENNEL, BULB; FRUITING VEGETABLES, CUCURBITS; GARLIC; LEAFY VEGETABLES; LEEK; ONION, BULB; PEAS (PODS AND SUCCULENT, IMMATURE SEEDS); POTATO; PULSES; SPINACH; SPRING ONION; TOMATO]	T7
CHLORPYRIFOS	
CHLORPYRIFOS	
CHERRIES	1
CRANBERRY	1
STONE FRUITS [EXCEPT CHERRIES]	T1
CLOTHIANIDIN	
CLOTHIANIDIN	
DRIED GRAPES	10
GRAPES [EXCEPT WINE GRAPES]	3
WINE GRAPES	*0.02
CYFLUTHRIN	
CYFLUTHRIN, SUM OF ISOMERS	
CHIA	T0.5
PAPAYA (PAWPAW)	T0.2
CYHALOTHRIN	
CYHALOTHRIN, SUM OF ISOMERS	
STONE FRUITS	0.5
CYPERMETHRIN	
CYPERMETHRIN, SUM OF ISOMERS	
DURIAN	1
LONGAN	1
PEPPERS, CHILI	1
EPOXICONAZOLE	
EPOXICONAZOLE	
CEREAL GRAINS	0.05
ETOXAZOLE	
ETOXAZOLE	
PODDED PEA (YOUNG PODS) (SNOW AND SUGAR SNAP)	T*0.02
FENBUCONAZOLE	
FENBUCONAZOLE	
WHEAT	*0.01

FENBUTATIN OXIDE	
BIS[TRIS(2-METHYL-2-PHENYLPROPYL)TIN]-OXIDE	
CHERRIES	6
FLUAZIFOP-BUTYL	
FLUAZIFOP-BUTYL	
CHIA	T2
ONION, WELSH	0.05
FLUBENDIAMIDE	
COMMODITIES OF PLANT ORIGIN: FLUBENDIAMIDE COMMODITIES OF ANIMAL ORIGIN: SUM OF FLUBENDIAMIDE AND 3-iodo-N-(2-methyl-4- [1,2,2,2-tetrafluoro-1- (trifluoromethyl)ethyl]phenyl)phthalimide, EXPRESSED AS FLUBENDIAMIDE	
EDIBLE OFFAL (MAMMALIAN)	0.03
MEAT (MAMMALIAN) (IN THE FAT)	0.05
MILK FATS	0.05
MILKS	*0.01
IMAZAMOX	
IMAZAMOX	
POPPY SEED	T*0.05
IMAZAPYR	
IMAZAPYR	
POPPY SEED	T*0.05
IMIDACLOPRID	
SUM OF IMIDACLOPRID AND METABOLITES CONTAINING THE 6-CHLOROPYRIDINYLMETHYLENE MOIETY, EXPRESSED AS IMIDACLOPRID	
BROAD BEAN (DRY)	*0.05
FIELD PEA (DRY)	*0.05
LEAFY VEGETABLES [EXCEPT LETTUCE, HEAD]	20
LENTIL (DRY)	0.2
LETTUCE, HEAD	5
INDOXACARB	
SUM OF INDOXACARB AND ITS R-ISOMER	
PEANUT	T0.02
IPRODIONE	
IPRODIONE	
PEPPERS	T2
METALAXYL	
METALAXYL	
GINGER, ROOT	T0.5
METHOMYL	
SUM OF METHOMYL AND METHYL HYDROXYTHIOACETIMIDATE ('METHOMYL OXIME'), EXPRESSED AS METHOMYL SEE ALSO THIODICARB	
CHIA	T0.5

METHOXYFENOZIDE	
METHOXYFENOZIDE	
CORIANDER (LEAVES, STEM, ROOTS)	T20
CRANBERRY	0.5
HERBS	T20
MEXICAN TARRAGON	T20
RUCOLA (ROCKET)	T20
STONE FRUITS [EXCEPT PLUMS (INCLUDING PRUNES)]	3
PACLOBUTRAZOL	
PACLOBUTRAZOL	
BARLEY	T0.1
WHEAT	T0.1
PENDIMETHALIN	
PENDIMETHALIN	
HERBS	*0.05
PERMETHRIN	
PERMETHRIN, SUM OF ISOMERS	
CORIANDER (LEAVES, STEM, ROOTS)	30
LEMON BALM	30
PHOSPHOROUS ACID	
PHOSPHOROUS ACID	
GINGER, ROOT	T100
TOMATO	T100
PIRIMICARB	
SUM OF PIRIMICARB, DEMETHYL-PIRIMICARB AND THE N-FORMYL-(METHYLAMINO) ANALOGUE (DEMETHYLFORMAMIDO-PIRIMICARB), EXPRESSED AS PIRIMICARB	
ADZUKI BEAN (DRY)	T0.5
LEAFY VEGETABLES [EXCEPT CHERVIL; MIZUNA; RUCOLA (ROCKET)]	T7
MUNG BEAN (DRY)	T0.5
ONION, WELSH	T3
SHALLOT	T3
SPRING ONION	T3
VEGETABLES [EXCEPT ADZUKI BEAN (DRY); LEAFY VEGETABLES; LUPIN (DRY); MUNG BEAN (DRY); ONION, WELSH; SHALLOT; SOYA BEAN (DRY); SPRING ONION; SWEET CORN (CORN-ON-THE- COB)]	1
PROFENOFOS	
PROFENOFOS	
MANGOSTEEN	5

PROTHIOCONAZOLE	
<i>COMMODITIES OF PLANT ORIGIN:</i> SUM OF PROTHIOCONAZOLE AND PROTHIOCONAZOLE DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLOROPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), EXPRESSED AS PROTHIOCONAZOLE	
<i>COMMODITIES OF ANIMAL ORIGIN:</i> SUM OF PROTHIOCONAZOLE, PROTHIOCONAZOLE DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLOROPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), PROTHIOCONAZOLE-3-HYDROXY-DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLORO-3-HYDROXYPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL) AND PROTHIOCONAZOLE-4-HYDROXY-DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLORO-4-HYDROXYPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), EXPRESSED AS PROTHIOCONAZOLE	
CEREAL BRAN, UNPROCESSED	0.5
OATS	*0.05
WHEAT GERM	0.5
PYRACLOSTROBIN	
<i>COMMODITIES OF PLANT ORIGIN:</i> PYRACLOSTROBIN	
<i>COMMODITIES OF ANIMAL ORIGIN:</i> SUM OF PYRACLOSTROBIN AND METABOLITES HYDROLYSED TO 1-(4-CHLORO-PHENYL)-1H-PYRAZOL-3-OL, EXPRESSED AS PYRACLOSTROBIN	
CEREAL GRAINS	*0.01
CUSTARD APPLE	T3
PAPAYA (PAWPAW)	T0.5
PYRIMETHANIL	
PYRIMETHANIL	
LEAFY VEGETABLES	T5
SPIROTETRAMAT	
SUM OF SPIROTETRAMAT, AND CIS-3-(2,5-DIMETHYLPHENYL)-4-HYDROXY-8-METHOXY-1-AZASPIRO[4.5]DEC-3-EN-2-ONE, EXPRESSED AS SPIROTETRAMAT	
DRIED GRAPES	4
FRUITING VEGETABLES, CUCURBITS [EXCEPT MELONS]	2
FRUITING VEGETABLES, OTHER THAN CUCURBITS	7
GRAPES	2
LEAFY VEGETABLES [EXCEPT LETTUCE, HEAD]	5
LEGUME VEGETABLES	T2
MELONS, EXCEPT WATERMELON	0.5
POTATO	5
SWEET POTATO	5
WATERMELON	0.5

TEBUCONAZOLE	
TEBUCONAZOLE	
CHERRIES	5
TEBUFENOZIDE	
TEBUFENOZIDE	
CRANBERRY	0.5
TERBUTHYLAZINE	
TERBUTHYLAZINE	
MAIZE	T*0.02
SORGHUM	T*0.02
SWEET CORN (CORN-ON-THE-COB)	T*0.02
TRIADIMENOL	
TRIADIMENOL SEE ALSO TRIADIMEFON	
PEPPERS	T1
TRICHLORFON	
TRICHLORFON	
FISH MUSCLE	T*0.01
FRUIT [EXCEPT BANANA; DRIED FRUITS; PEACH]	0.1
VEGETABLES [EXCEPT BEETROOT; BRUSSELS SPROUTS; CAULIFLOWER; CELERY; KALE; PEPPERS; PULSES; SUGAR BEET; SWEET CORN (CORN-ON-THE-COB)]	0.1
TRIFLOXYSTROBIN	
SUM OF TRIFLOXYSTROBIN AND ITS ACID METABOLITE ((E,E)-METHOXYIMINO-[2-[1-(3-TRIFLUOROMETHYLPHENYL)-ETHYLIDENEAMINOXYMETHYL]PHENYL] ACETIC ACID), EXPRESSED AS TRIFLOXYSTROBIN EQUIVALENTS	
CELERY	T1
CHARD (SILVER BEET)	T0.7
CHICORY LEAVES	T0.7
ENDIVE	T0.7
SPINACH	T0.7
STONE FRUITS	2
TRIFLURALIN	
TRIFLURALIN	
CHIA	T*0.01
TRINEXAPAC-ETHYL	
4-(CYCLOPROPYL- α -HYDROXY-METHYLENE)-3,5-DIOXO-CYCLOHEXANECARBOXYLIC ACID	
BARLEY	T0.3
WHEAT	T0.3

[1.5] omitting from Schedule 1, under the entries for the following chemicals, the Maximum Residue Limit for the food, substituting –

BIFENTHRIN BIFENTHRIN	
PEAS (PODS AND SUCCULENT, IMMATURE SEEDS)	*0.01
BOSCALID COMMODITIES OF PLANT ORIGIN: BOSCALID COMMODITIES OF ANIMAL ORIGIN: SUM OF BOSCALID, 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE AND THE GLUCURONIDE CONJUGATE OF 2-CHLORO-N-(4'-CHLORO-5-HYDROXYBIPHENYL-2-YL) NICOTINAMIDE, EXPRESSED AS BOSCALID EQUIVALENTS	
BRASSICA LEAFY VEGETABLES	T30
EDIBLE OFFAL (MAMMALIAN)	0.3
LETTUCE, HEAD	T15
LETTUCE, LEAF	T15
MEAT (MAMMALIAN) (IN THE FAT)	0.3
MILKS	0.1
BROMOXYNIL BROMOXYNIL	
EDIBLE OFFAL (MAMMALIAN)	T3
MEAT (MAMMALIAN) (IN THE FAT)	T1
MILKS	T0.1
CHLOROTHALONIL COMMODITIES OF PLANT ORIGIN: CHLOROTHALONIL COMMODITIES OF ANIMAL ORIGIN: SUM OF CHLOROTHALONIL AND 4-HYDROXY-2, 5, 6-TRICHLOROISOPHTHALONITRILE METABOLITE, EXPRESSED AS CHLOROTHALONIL	
EDIBLE OFFAL (MAMMALIAN)	7
HERBS [EXCEPT FENNEL, LEAF]	T20
MEAT (MAMMALIAN) (IN THE FAT)	2
MILKS	0.05
PULSES	3
CHLORPYRIFOS CHLORPYRIFOS	
BLUEBERRIES	*0.01
FENBUCONAZOLE FENBUCONAZOLE	
EDIBLE OFFAL (MAMMALIAN)	0.05
STONE FRUITS [EXCEPT NECTARINE]	1

FIPRONIL SUM OF FIPRONIL, THE SULPHENYL METABOLITE (5-AMINO-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-4-[(TRIFLUOROMETHYL)SULPHENYL]-1H-PYRAZOLE-3-CARBONITRILE), THE SULPHONYL METABOLITE (5-AMINO-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-4-[(TRIFLUOROMETHYL)SULPHONYL]-1H-PYRAZOLE-3-CARBONITRILE), AND THE TRIFLUOROMETHYL METABOLITE (5-AMINO-4-TRIFLUOROMETHYL-1-[2,6-DICHLORO-4-(TRIFLUOROMETHYL)PHENYL]-1H-PYRAZOLE-3-CARBONITRILE)	
SWEET POTATO	*0.01
FLUAZIFOP-BUTYL FLUAZIFOP-BUTYL	
PARSNIP	0.1
FLUBENDIAMIDE COMMODITIES OF PLANT ORIGIN: FLUBENDIAMIDE COMMODITIES OF ANIMAL ORIGIN: SUM OF FLUBENDIAMIDE AND 3-iodo-N-(2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl)phthalimide, EXPRESSED AS FLUBENDIAMIDE	
LETTUCE, HEAD	5
LETTUCE, LEAF	7
PEPPERS, SWEET	1
TOMATO	2
FLUMETSULAM FLUMETSULAM	
EDIBLE OFFAL (MAMMALIAN)	0.3
IMIDACLOPRID SUM OF IMIDACLOPRID AND METABOLITES CONTAINING THE 6-CHLOROPYRIDINYL METHYLENE MOIETY, EXPRESSED AS IMIDACLOPRID	
LUPIN (DRY)	0.2
POTATO	0.3
SWEET POTATO	0.3
IPRODIONE IPRODIONE	
BRUSSELS SPROUTS	0.5
METALAXYL METALAXYL	
PAPAYA (PAWPAW)	*0.01
PERMETHRIN PERMETHRIN, SUM OF ISOMERS	
HERBS	30
KAFFIR LIME LEAVES	30

LEMON GRASS	30
PROTHIOCONAZOLE	
COMMODITIES OF PLANT ORIGIN: SUM OF PROTHIOCONAZOLE AND PROTHIOCONAZOLE DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLOROPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), EXPRESSED AS PROTHIOCONAZOLE	
COMMODITIES OF ANIMAL ORIGIN: SUM OF PROTHIOCONAZOLE, PROTHIOCONAZOLE DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLOROPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), PROTHIOCONAZOLE-3-HYDROXY-DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLORO-3-HYDROXYPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL) AND PROTHIOCONAZOLE-4-HYDROXY-DESTHIO (2-(1-CHLOROCYCLOPROPYL)-1-(2-CHLORO-4-HYDROXYPHENYL)-3-(1H-1,2,4-TRIAZOL-1-YL)-PROPAN-2-OL), EXPRESSED AS PROTHIOCONAZOLE	
BARLEY	0.3
EDIBLE OFFAL (MAMMALIAN)	0.1
WHEAT	0.3
PYRIPROXYFEN	
PYRIPROXYFEN	
MANGO	0.05
SIMAZINE	
SIMAZINE	
EDIBLE OFFAL (MAMMALIAN)	*0.05


MEAT (MAMMALIAN)	*0.05
MILKS	*0.02
SPIROTETRAMAT	
SUM OF SPIROTETRAMAT, AND CIS-3-(2,5-DIMETHYLPHENYL)-4-HYDROXY-8-METHOXY-1-AZASPIRO[4.5]DEC-3-EN-2-ONE, EXPRESSED AS SPIROTETRAMAT	
CITRUS FRUITS	1
LETTUCE, HEAD	3
MANGO	0.3
ONION, BULB	0.5
TERBUTHYLAZINE	
TERBUTHYLAZINE	
EDIBLE OFFAL (MAMMALIAN)	*0.01
EGGS	*0.01
MEAT (MAMMALIAN)	*0.01
MILKS	*0.01
POULTRY, EDIBLE OFFAL OF	*0.01
POULTRY MEAT	*0.01
PULSES	*0.02
RAPE SEED (CANOLA)	*0.02
TOLCLOFOS-METHYL	
TOLCLOFOS-METHYL	
BEETROOT	*0.01

Summary of proposed MRLs and technical amendments in Proposal M1006

INTERPRETIVE GUIDE TO THE SUMMARY TABLE OF MRLS

The following is an example of an entry and the proposed MRL is not being considered in this Proposal. Further information on calculating dietary exposure is provided at [Supporting Document 1](#).

Data from the 19th and 20th ATDS are provided when available because they provide an indication of the typical exposure to chemicals in table ready foods. The ATDS results are more realistic because analysed concentrations of the chemical in foods as consumed are used. The National Estimated Daily Intake (NEDI) and National Estimated Short Term Intake (NESTI) calculations are theoretical calculations that protectively overestimate exposure. Small variations may be noted in the exposure assessment between different ATDSs. These variations are minor and are typically due to the different range of foods in the individual studies.



**SUMMARY OF MRLS UNDER CONSIDERATION IN PROPOSAL M1006
APVMA MRLS – OCTOBER 2009 – MARCH 2010 AND OTHER REQUESTS**

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment						
<p>Abamectin Abamectin is an insecticide and acaricide with contact and stomach action. It inhibits stimulation of neurons by binding to gamma-aminobutyric acid regulated chloride channels and allowing free passage of chloride ions into the neuron. It is used to control mites on cotton and various fruits and vegetables.</p> <p>The APVMA has issued a permit for its use to control two-spotted mite (<i>Tetranychus urticae</i>) on sweet corn. The recommended temporary MRL is at the limit of analytical quantification (LOQ).</p> <p>Sweet corn (corn-on-the-cob) Insert T*0.01</p>	<p>NEDI: 89% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0"> <tr> <td align="center"><u>2-6 years</u></td> <td align="center"><u>2+ years</u></td> </tr> <tr> <td align="center">6</td> <td align="center">2</td> </tr> </table>	<u>2-6 years</u>	<u>2+ years</u>	6	2		
<u>2-6 years</u>	<u>2+ years</u>						
6	2						
<p>Benzyladenine Benzyladenine is a plant growth regulator. It stimulates protein synthesis. It is a synthetic cytokinin. Its limited translocation is utilised to restrict effects to the target part of the plant. It is used to regulate bud emergence and fruit set, increase fruit size and stimulate flower bud formation and regular bearing in fruit trees.</p> <p>The APVMA has issued a permit for its use to reduce alternate bearing in pistachios. The recommended temporary MRL is at the LOQ.</p> <p>Pistachio nut Insert T*0.05</p>	<p>NEDI: 1% of the ADI</p>						
<p>Beta-cyfluthrin Beta cyfluthrin is a non-systemic pyrethroid insecticide with contact and stomach action. It acts on the nervous system of insects and disturbs the function of neurons by interaction with the sodium channel. It is used to control a range of pests including Lepidoptera and Homoptera on many crops.</p> <p>The APVMA has issued permits for its use to control heliothis in chia and fruit-spotting bug (<i>Amblypelta nitida</i>) and banana-spotting bug (<i>Amblypelta lutescens lutescens</i>) in pawpaw.</p> <p>Note: Beta-cyfluthrin MRLs are listed under cyfluthrin.</p> <p>Chia Insert T0.5 Papaya (pawpaw) Insert T0.2</p>	<p>NEDI: 68% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0"> <tr> <td align="center"><u>2-6 years</u></td> <td align="center"><u>2+ years</u></td> </tr> <tr> <td align="center"><1</td> <td align="center"><1</td> </tr> <tr> <td align="center">18</td> <td align="center">5</td> </tr> </table>	<u>2-6 years</u>	<u>2+ years</u>	<1	<1	18	5
<u>2-6 years</u>	<u>2+ years</u>						
<1	<1						
18	5						

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																	
<p>Bifenazate Bifenazate is a non-systemic acaricide. It is a neuronal inhibitor with predominantly contact action and long residual action. It is used to control the egg and motile stages of phytophagous mites on various crops.</p> <p>The APVMA has issued a permit for its use to control two-spotted mite (<i>Tetranychus urticae</i>) on leafy and head lettuce varieties grown in protected situations.</p> <p>The United States Northwest Horticultural Council (NHC) requested that FSANZ include an MRL in the Code harmonised with the United States limit for bifenazate residues in cherries. Bifenazate residues may occur in cherries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <table border="0" data-bbox="177 772 983 869"> <tr> <td>Cherries</td> <td>Insert</td> <td>2.5</td> </tr> <tr> <td>Lettuce, head</td> <td>Insert</td> <td>T5</td> </tr> <tr> <td>Lettuce, leaf</td> <td>Insert</td> <td>T5</td> </tr> </table>	Cherries	Insert	2.5	Lettuce, head	Insert	T5	Lettuce, leaf	Insert	T5	<p>NEDI: 24% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 739 1388 869"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td></td> <td>14</td> <td>12</td> </tr> <tr> <td></td> <td>4</td> <td>2</td> </tr> <tr> <td></td> <td>1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>		14	12		4	2		1	<1												
Cherries	Insert	2.5																																
Lettuce, head	Insert	T5																																
Lettuce, leaf	Insert	T5																																
	<u>2-6 years</u>	<u>2+ years</u>																																
	14	12																																
	4	2																																
	1	<1																																
<p>Bifenthrin Bifenthrin is a synthetic pyrethroid insecticide. It kills insects by affecting the salt balance (sodium channels) in nerve cells. It has a broad spectrum of activity against insects with the main toxic effect on the nervous system. It is used to control a broad range of foliar pests on cereal, fruit and vegetable crops.</p> <p>The APVMA has issued permits for its use to control symphylids, ground dwelling insects, (<i>Hanseniella</i> spp.) in pineapple, silverleaf whitefly on cucumbers and red-legged earth mite and blue oat mite in peas. The recommended MRLs for pineapple and peas are at the LOQ.</p> <table border="0" data-bbox="177 1265 983 1505"> <tr> <td>Cucumber</td> <td>Insert</td> <td>T0.3</td> </tr> <tr> <td>Fruiting vegetables, cucurbits</td> <td>Omit</td> <td>0.1</td> </tr> <tr> <td>Fruiting vegetables, cucurbits [except cucumber]</td> <td>Insert</td> <td>0.1</td> </tr> <tr> <td>Peas (pods and succulent, immature seeds)</td> <td>Omit</td> <td>T*0.01</td> </tr> <tr> <td></td> <td>Substitute</td> <td>*0.01</td> </tr> <tr> <td>Pineapple</td> <td>Insert</td> <td>T*0.01</td> </tr> </table>	Cucumber	Insert	T0.3	Fruiting vegetables, cucurbits	Omit	0.1	Fruiting vegetables, cucurbits [except cucumber]	Insert	0.1	Peas (pods and succulent, immature seeds)	Omit	T*0.01		Substitute	*0.01	Pineapple	Insert	T*0.01	<p>NEDI: 76% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1220 1388 1505"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td></td> <td>48</td> <td>17</td> </tr> <tr> <td></td> <td>60</td> <td>16</td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td></td> <td>10</td> <td>3</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>		48	17		60	16		<1	<1		10	3
Cucumber	Insert	T0.3																																
Fruiting vegetables, cucurbits	Omit	0.1																																
Fruiting vegetables, cucurbits [except cucumber]	Insert	0.1																																
Peas (pods and succulent, immature seeds)	Omit	T*0.01																																
	Substitute	*0.01																																
Pineapple	Insert	T*0.01																																
	<u>2-6 years</u>	<u>2+ years</u>																																
	48	17																																
	60	16																																
	<1	<1																																
	10	3																																

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																																																																																															
<p>Chlorothalonil Chlorothalonil is a non-systemic foliar fungicide with protective action. It conjugates with and depletes thiols, particularly glutathione, in germinating fungal cells leading to disruption of glycolysis and energy production. It is used to control fungal diseases in a broad range of crops and horticultural situations.</p> <p>The APVMA has approved an extension of use of chlorothalonil to include chickpeas and lentils and a use pattern for silverbeet and spinach. The APVMA has also issued permits for its use to control downy mildew, Alternaria, Botrytis and Cercospora on certain culinary herbs; and with pyrimethanil to control Alternaria and Botrytis on chickory, endive, radicchio, silverbeet and spinach.</p> <p>Amendment to residue definition</p> <p>Omit: Commodities of plant origin: Chlorothalonil Commodities of animal origin: Sum of chlorothalonil and 4-hydroxy-2,5,6-trichloroisophthalonitrile metabolite, expressed as chlorothalonil</p> <p>Substitute: Commodities of plant origin: Chlorothalonil Commodities of animal origin: 4-hydroxy-2,5,6-trichloroisophthalonitrile metabolite, expressed as chlorothalonil</p> <table border="0" data-bbox="177 1048 983 1930"> <tr> <td>Chard (silver beet)</td> <td>Insert</td> <td>T50</td> </tr> <tr> <td>Coriander (leaves, stem, roots)</td> <td>Insert</td> <td>T20</td> </tr> <tr> <td>Edible offal (mammalian)</td> <td>Omit</td> <td>T3</td> </tr> <tr> <td></td> <td>Substitute</td> <td>7</td> </tr> <tr> <td>Herbs [except fennel, leaf]</td> <td>Omit</td> <td>T7</td> </tr> <tr> <td></td> <td>Substitute</td> <td>T20</td> </tr> <tr> <td>Leafy vegetables</td> <td>Omit</td> <td>T7</td> </tr> <tr> <td>Leafy vegetables [except chard (silver beet); spinach]</td> <td>Insert</td> <td>T10</td> </tr> <tr> <td>Meat (mammalian) (in the fat)</td> <td>Omit</td> <td>T2</td> </tr> <tr> <td></td> <td>Substitute</td> <td>2</td> </tr> <tr> <td>Milks</td> <td>Omit</td> <td>T0.05</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.05</td> </tr> <tr> <td>Poultry, edible offal of</td> <td>Insert</td> <td>*0.05</td> </tr> <tr> <td>Poultry meat</td> <td>Insert</td> <td>*0.05</td> </tr> <tr> <td>Pulses</td> <td>Omit</td> <td>T7</td> </tr> <tr> <td></td> <td>Substitute</td> <td>3</td> </tr> <tr> <td>Spinach</td> <td>Insert</td> <td>T100</td> </tr> <tr> <td>Vegetables [except as otherwise listed under this chemical]</td> <td>Omit</td> <td>T7</td> </tr> <tr> <td>Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]</td> <td>Insert</td> <td>T7</td> </tr> </table>	Chard (silver beet)	Insert	T50	Coriander (leaves, stem, roots)	Insert	T20	Edible offal (mammalian)	Omit	T3		Substitute	7	Herbs [except fennel, leaf]	Omit	T7		Substitute	T20	Leafy vegetables	Omit	T7	Leafy vegetables [except chard (silver beet); spinach]	Insert	T10	Meat (mammalian) (in the fat)	Omit	T2		Substitute	2	Milks	Omit	T0.05		Substitute	0.05	Poultry, edible offal of	Insert	*0.05	Poultry meat	Insert	*0.05	Pulses	Omit	T7		Substitute	3	Spinach	Insert	T100	Vegetables [except as otherwise listed under this chemical]	Omit	T7	Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]	Insert	T7	<p>NEDI: 88% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>19th ATDS: <1% of the ADI for all population groups assessed</p> <p>Note that the proposed vegetables MRL variation is a technical amendment only. NESTI calculations are not required.</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1016 1390 1601"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td>Chard (silver beet)</td> <td>19</td> <td>11</td> </tr> <tr> <td>Coriander (leaves, stem, roots)</td> <td>2</td> <td>1</td> </tr> <tr> <td>Edible offal (mammalian)</td> <td></td> <td></td> </tr> <tr> <td>Herbs [except fennel, leaf]</td> <td>1</td> <td>2</td> </tr> <tr> <td>Leafy vegetables</td> <td>2</td> <td>1</td> </tr> <tr> <td>Leafy vegetables [except chard (silver beet); spinach]</td> <td>15</td> <td>8</td> </tr> <tr> <td>Meat (mammalian) (in the fat)</td> <td></td> <td></td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td>Milks</td> <td></td> <td></td> </tr> <tr> <td></td> <td>1</td> <td>1</td> </tr> <tr> <td>Poultry, edible offal of</td> <td></td> <td></td> </tr> <tr> <td>Poultry meat</td> <td><1</td> <td><1</td> </tr> <tr> <td>Pulses</td> <td></td> <td></td> </tr> <tr> <td></td> <td>3</td> <td>1</td> </tr> <tr> <td>Spinach</td> <td>38</td> <td>54</td> </tr> <tr> <td>Vegetables [except as otherwise listed under this chemical]</td> <td></td> <td></td> </tr> <tr> <td>Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]</td> <td></td> <td></td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	Chard (silver beet)	19	11	Coriander (leaves, stem, roots)	2	1	Edible offal (mammalian)			Herbs [except fennel, leaf]	1	2	Leafy vegetables	2	1	Leafy vegetables [except chard (silver beet); spinach]	15	8	Meat (mammalian) (in the fat)				<1	<1	Milks				1	1	Poultry, edible offal of			Poultry meat	<1	<1	Pulses				3	1	Spinach	38	54	Vegetables [except as otherwise listed under this chemical]			Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]		
Chard (silver beet)	Insert	T50																																																																																																														
Coriander (leaves, stem, roots)	Insert	T20																																																																																																														
Edible offal (mammalian)	Omit	T3																																																																																																														
	Substitute	7																																																																																																														
Herbs [except fennel, leaf]	Omit	T7																																																																																																														
	Substitute	T20																																																																																																														
Leafy vegetables	Omit	T7																																																																																																														
Leafy vegetables [except chard (silver beet); spinach]	Insert	T10																																																																																																														
Meat (mammalian) (in the fat)	Omit	T2																																																																																																														
	Substitute	2																																																																																																														
Milks	Omit	T0.05																																																																																																														
	Substitute	0.05																																																																																																														
Poultry, edible offal of	Insert	*0.05																																																																																																														
Poultry meat	Insert	*0.05																																																																																																														
Pulses	Omit	T7																																																																																																														
	Substitute	3																																																																																																														
Spinach	Insert	T100																																																																																																														
Vegetables [except as otherwise listed under this chemical]	Omit	T7																																																																																																														
Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]	Insert	T7																																																																																																														
	<u>2-6 years</u>	<u>2+ years</u>																																																																																																														
Chard (silver beet)	19	11																																																																																																														
Coriander (leaves, stem, roots)	2	1																																																																																																														
Edible offal (mammalian)																																																																																																																
Herbs [except fennel, leaf]	1	2																																																																																																														
Leafy vegetables	2	1																																																																																																														
Leafy vegetables [except chard (silver beet); spinach]	15	8																																																																																																														
Meat (mammalian) (in the fat)																																																																																																																
	<1	<1																																																																																																														
Milks																																																																																																																
	1	1																																																																																																														
Poultry, edible offal of																																																																																																																
Poultry meat	<1	<1																																																																																																														
Pulses																																																																																																																
	3	1																																																																																																														
Spinach	38	54																																																																																																														
Vegetables [except as otherwise listed under this chemical]																																																																																																																
Vegetables [except asparagus; Brussels sprouts; carrot; celery; chard (silver beet); fennel, bulb; fruiting vegetables, cucurbits; garlic; leafy vegetables; leek; onion, bulb; peas (pods and succulent, immature seeds); potato; pulses; spinach; spring onion; tomato]																																																																																																																

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																														
<p>Chlorpyrifos Chlorpyrifos is a broad spectrum non-systemic insecticide with contact, stomach, and respiratory action. It is a cholinesterase inhibitor. It is used to control a broad range of insect pests in many crops including cotton, sugarcane, vegetables, pome and stone fruit, pastures, turf and ornamental crops.</p> <p>The APVMA has evaluated trial data in relation to an existing permit to use chlorpyrifos to control scarab beetles (<i>Scarabaeidae</i>) on blueberries. The recommended MRL is at the LOQ.</p> <p>The NHC requested that FSANZ consider including a chlorpyrifos MRL in the Code harmonised with the United States MRL for chlorpyrifos residues in cherries.</p> <p>The CMC requested that FSANZ include an MRL in the Code harmonised with the Codex limit for chlorpyrifos residues in cranberries.</p> <p>Chlorpyrifos residues may occur in cherries and cranberries imported from the United States. The proposed MRLs may minimise potential trade disruption and extend consumer choice.</p> <p>Chlorpyrifos is currently under review by the APVMA. FSANZ notes that the conclusion of the review is imminent and that upon finalisation, the APVMA may vary chlorpyrifos MRLs. Following the anticipated recommended changes to use patterns, the estimated dietary exposures will be reassessed as part of finalisation of the Review. Further information about the review is available on the APVMA website at: www.apvma.gov.au/products/review/current/chlorpyrifos.php</p> <table border="0" data-bbox="177 1263 983 1447"> <tr> <td>Blueberries</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td></td> <td>Substitute</td> <td>*0.01</td> </tr> <tr> <td>Cherries</td> <td>Insert</td> <td>1</td> </tr> <tr> <td>Cranberry</td> <td>Insert</td> <td>1</td> </tr> <tr> <td>Stone fruits</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td>Stone fruits [except cherries]</td> <td>Insert</td> <td>T1</td> </tr> </table>	Blueberries	Omit	T1		Substitute	*0.01	Cherries	Insert	1	Cranberry	Insert	1	Stone fruits	Omit	T1	Stone fruits [except cherries]	Insert	T1	<p>NEDI: 75% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>19th ATDS: 3% of the ADI for toddlers 2 years; 1% of the ADI for boys 12 years and <1% of the ADI for other population groups assessed</p> <p>Note that the proposed stone fruits MRL variation is a technical amendment only. NESTI calculations are not required.</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1211 1390 1384"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td><1</td> <td><1</td> <td><1</td> </tr> <tr> <td>16</td> <td>3</td> <td>3</td> </tr> <tr> <td><1</td> <td><1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	<1	<1	<1	16	3	3	<1	<1	<1
Blueberries	Omit	T1																													
	Substitute	*0.01																													
Cherries	Insert	1																													
Cranberry	Insert	1																													
Stone fruits	Omit	T1																													
Stone fruits [except cherries]	Insert	T1																													
	<u>2-6 years</u>	<u>2+ years</u>																													
<1	<1	<1																													
16	3	3																													
<1	<1	<1																													
<p>Clothianidin Clothianidin is an insecticide. It is an agonist of the nicotinic acetylcholine receptor, affecting the synapses in the insect central nervous system. It exhibits translaminar and root systemic activity. It is used to control various pests in pome and stone fruits, bananas and cotton.</p> <p>The APVMA has approved an extension of its use to control long tail mealybug in grapes. The recommended MRL for wine grapes is at the LOQ.</p> <table border="0" data-bbox="177 1787 983 1872"> <tr> <td>Dried grapes</td> <td>Insert</td> <td>10</td> </tr> <tr> <td>Grapes [except wine grapes]</td> <td>Insert</td> <td>3</td> </tr> <tr> <td>Wine grapes</td> <td>Insert</td> <td>*0.02</td> </tr> </table>	Dried grapes	Insert	10	Grapes [except wine grapes]	Insert	3	Wine grapes	Insert	*0.02	<p>NEDI: 4% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1727 1390 1872"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td>12</td> <td>3</td> <td>3</td> </tr> <tr> <td>27</td> <td>11</td> <td>11</td> </tr> <tr> <td><1</td> <td><1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	12	3	3	27	11	11	<1	<1	<1									
Dried grapes	Insert	10																													
Grapes [except wine grapes]	Insert	3																													
Wine grapes	Insert	*0.02																													
	<u>2-6 years</u>	<u>2+ years</u>																													
12	3	3																													
27	11	11																													
<1	<1	<1																													

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																	
<p>Cypermethrin Cypermethrin is a pyrethroid, non-systemic insecticide with contact and stomach action. It acts on the central and peripheral nervous system of insects in very low doses. It is used to control a wide range of chewing and sucking insect pests in cereal, legume and oilseed crops and horticultural situations.</p> <p>The Thailand National Bureau of Agricultural Commodity and Food Standards has requested that FSANZ consider including MRLs in the Code harmonised with the Thai MRLs for cypermethrin residues in durians, longans and chillies. Cypermethrin residues may occur in imported fruits. The proposed MRLs may minimise potential trade disruption and extend consumer choice.</p> <p>The commodity name 'Peppers, Chili' is used for chillies consistent with the Codex classification of foods and animal feeds.</p> <table border="0" data-bbox="177 831 983 987"> <tr> <td>Durian</td> <td>Insert</td> <td>1</td> </tr> <tr> <td>Longan</td> <td>Insert</td> <td>1</td> </tr> <tr> <td>Peppers, Chili</td> <td>Insert</td> <td>1</td> </tr> </table>	Durian	Insert	1	Longan	Insert	1	Peppers, Chili	Insert	1	<p>NEDI: 12% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: not detected in any foods sampled</p> <p>19th ATDS: <1% of the ADI for all population groups assessed</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 775 1390 987"> <tr> <td></td> <td><u>2-6 years</u></td> <td></td> <td><u>2+ years</u></td> </tr> <tr> <td>79</td> <td>Tropical fruit</td> <td>20</td> <td></td> </tr> <tr> <td></td> <td>inedible peel</td> <td></td> <td></td> </tr> <tr> <td>79</td> <td>Tropical fruit</td> <td>20</td> <td></td> </tr> <tr> <td></td> <td>inedible peel</td> <td></td> <td></td> </tr> <tr> <td>24</td> <td>Peppers group</td> <td>10</td> <td></td> </tr> </table>		<u>2-6 years</u>		<u>2+ years</u>	79	Tropical fruit	20			inedible peel			79	Tropical fruit	20			inedible peel			24	Peppers group	10	
Durian	Insert	1																																
Longan	Insert	1																																
Peppers, Chili	Insert	1																																
	<u>2-6 years</u>		<u>2+ years</u>																															
79	Tropical fruit	20																																
	inedible peel																																	
79	Tropical fruit	20																																
	inedible peel																																	
24	Peppers group	10																																
<p>Epoxiconazole Epoxiconazole is a broad spectrum contact and systemic fungicide. It inhibits C-14 demethylase in sterol biosynthesis. It is used to control various fungal diseases in a range of crops and horticultural situations.</p> <p>The APVMA has approved its use with pyraclostrobin to control various diseases in wheat, barley and oats.</p> <table border="0" data-bbox="177 1267 983 1357"> <tr> <td>Barley</td> <td>Omit</td> <td>0.05</td> </tr> <tr> <td>Cereal grains</td> <td>Insert</td> <td>0.05</td> </tr> <tr> <td>Wheat</td> <td>Omit</td> <td>0.05</td> </tr> </table>	Barley	Omit	0.05	Cereal grains	Insert	0.05	Wheat	Omit	0.05	<p>NEDI: 2% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1211 1390 1357"> <tr> <td></td> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td><1</td> <td></td> <td><1</td> </tr> </table>		<u>2-6 years</u>	<u>2+ years</u>	<1		<1																		
Barley	Omit	0.05																																
Cereal grains	Insert	0.05																																
Wheat	Omit	0.05																																
	<u>2-6 years</u>	<u>2+ years</u>																																
<1		<1																																
<p>Etoxazole Etoxazole is a contact acaricide and insect growth regulator. It inhibits the moulting process of mites and aphids by disrupting the cell wall. It is used to control various mites on pome fruit, stone fruit, table grapes and cotton.</p> <p>The APVMA has issued a permit for its use to control two-spotted mite (<i>Tetranychus urticae</i>) on snow peas and sugar snap peas. The recommended MRL is at the LOQ.</p> <table border="0" data-bbox="177 1659 983 1727"> <tr> <td>Podded pea (young pods) (snow and sugar snap)</td> <td>Insert</td> <td>T*0.02</td> </tr> </table>	Podded pea (young pods) (snow and sugar snap)	Insert	T*0.02	<p>NEDI: 2% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1626 1390 1727"> <tr> <td></td> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td><1</td> <td></td> <td><1</td> </tr> </table>		<u>2-6 years</u>	<u>2+ years</u>	<1		<1																								
Podded pea (young pods) (snow and sugar snap)	Insert	T*0.02																																
	<u>2-6 years</u>	<u>2+ years</u>																																
<1		<1																																

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																											
<p>Fenbuconazole Fenbuconazole is a systemic fungicide with protectant, curative and eradicant properties. It inhibits steroid demethylation. It is used to control certain diseases in bananas, nectarines and wheat.</p> <p>The APVMA has issued a permit for its use to control fungal disease in wheat. The recommended MRL is at the LOQ. An increased offal MRL is recommended as residues may occur in liver.</p> <p>The NHC requested that FSANZ consider including a fenbuconazole MRL in the Code harmonised with the United States MRL for fenbuconazole residues in cherries. Residues may occur in cherries imported from the United States. The proposed MRL may minimise potential trade disruption and extend consumer choice.</p> <table border="0" data-bbox="177 801 983 958"> <tr> <td>Edible offal (mammalian)</td> <td>Omit</td> <td>*0.01</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.05</td> </tr> <tr> <td>Stone fruits [except nectarine]</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td></td> <td>Substitute</td> <td>1</td> </tr> <tr> <td>Wheat</td> <td>Insert</td> <td>*0.01</td> </tr> </table>	Edible offal (mammalian)	Omit	*0.01		Substitute	0.05	Stone fruits [except nectarine]	Omit	T1		Substitute	1	Wheat	Insert	*0.01	<p>NEDI: 3% of the ADI</p> <p>Note that the proposed stone fruits MRL variation is a technical amendment only. NESTI calculations are not required.</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 741 1390 958"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td>Edible offal (mammalian)</td> <td><1</td> <td><1</td> </tr> <tr> <td>Stone fruits [except nectarine]</td> <td><1</td> <td><1</td> </tr> <tr> <td>Wheat</td> <td><1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	Edible offal (mammalian)	<1	<1	Stone fruits [except nectarine]	<1	<1	Wheat	<1	<1
Edible offal (mammalian)	Omit	*0.01																										
	Substitute	0.05																										
Stone fruits [except nectarine]	Omit	T1																										
	Substitute	1																										
Wheat	Insert	*0.01																										
	<u>2-6 years</u>	<u>2+ years</u>																										
Edible offal (mammalian)	<1	<1																										
Stone fruits [except nectarine]	<1	<1																										
Wheat	<1	<1																										
<p>Fenbutatin oxide Fenbutatin oxide is a non-systemic acaricide with contact and stomach action. It inhibits oxidative phosphorylation. It is used to control phytophagous mites in various horticultural situations.</p> <p>The NHC requested that FSANZ include an MRL in the Code harmonised with the United States limit for fenbutatin oxide residues in cherries. Residues may occur in cherries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <table border="0" data-bbox="177 1294 983 1328"> <tr> <td>Cherries</td> <td>Insert</td> <td>6</td> </tr> </table>	Cherries	Insert	6	<p>NEDI: 82% of the ADI</p>																								
Cherries	Insert	6																										
<p>Fipronil Fipronil is a phenylpyrazole insecticide. It blocks the GABA regulated chloride channel. This disrupts central nervous system activity. It is used to control pests in a wide range of crops and horticultural situations.</p> <p>The APVMA has approved a use pattern to control various pests on sweet potatoes. Residues data indicate that detectable residues are not expected to occur. The data are sufficient to remove the temporary status of the MRL.</p> <table border="0" data-bbox="177 1664 983 1722"> <tr> <td>Sweet potato</td> <td>Omit</td> <td>T*0.01</td> </tr> <tr> <td></td> <td>Substitute</td> <td>*0.01</td> </tr> </table>	Sweet potato	Omit	T*0.01		Substitute	*0.01	<p>NEDI: 77% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1603 1390 1722"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td>Sweet potato</td> <td>2</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	Sweet potato	2	<1															
Sweet potato	Omit	T*0.01																										
	Substitute	*0.01																										
	<u>2-6 years</u>	<u>2+ years</u>																										
Sweet potato	2	<1																										

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																				
<p>Fluazifop-butyl Fluazifop-butyl (fluazifop) is a selective systemic herbicide absorbed by the leaves. It inhibits acetyl-coA carboxylase. It is used to control grass weeds in broad leaf crops.</p> <p>The APVMA has issued permits for its use to control certain grass weeds in chia, parsnip and various onions.</p> <table border="0" data-bbox="177 501 983 622"> <tr> <td>Chia</td> <td>Insert</td> <td>T2</td> </tr> <tr> <td>Onion, Welsh</td> <td>Insert</td> <td>0.05</td> </tr> <tr> <td>Parsnip</td> <td>Omit</td> <td>T0.1</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.1</td> </tr> </table>	Chia	Insert	T2	Onion, Welsh	Insert	0.05	Parsnip	Omit	T0.1		Substitute	0.1	<p>NEDI: 69% of the ADI</p>																								
Chia	Insert	T2																																			
Onion, Welsh	Insert	0.05																																			
Parsnip	Omit	T0.1																																			
	Substitute	0.1																																			
<p>Flubendiamide Flubendiamide is an insecticide. It has larvicidal activity, when orally ingested it results in rapid cessation of feeding. It is a ryanodine receptor agonist. It is used to control insect pests in various vegetables.</p> <p>The APVMA has evaluated further data in relation to the approved use of flubendiamide in certain vegetables. The data are sufficient to remove the temporary status of the recommended vegetable MRLs. Residues may occur in certain stock feed following the approved use of flubendiamide. Animal commodity MRLs are recommended as data indicate residues may occur in these foods. The recommended milk MRL is at the LOQ.</p> <table border="0" data-bbox="177 1084 983 1451"> <tr> <td>Edible offal (mammalian)</td> <td>Insert</td> <td>0.03</td> </tr> <tr> <td>Lettuce, head</td> <td>Omit</td> <td>T5</td> </tr> <tr> <td></td> <td>Substitute</td> <td>5</td> </tr> <tr> <td>Lettuce, leaf</td> <td>Omit</td> <td>T5</td> </tr> <tr> <td></td> <td>Substitute</td> <td>7</td> </tr> <tr> <td>Meat (mammalian) (in the fat)</td> <td>Insert</td> <td>0.05</td> </tr> <tr> <td>Milk fats</td> <td>Insert</td> <td>0.05</td> </tr> <tr> <td>Milks</td> <td>Insert</td> <td>*0.01</td> </tr> <tr> <td>Peppers, Sweet</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td></td> <td>Substitute</td> <td>1</td> </tr> <tr> <td>Tomato</td> <td>Omit</td> <td>T2</td> </tr> <tr> <td></td> <td>Substitute</td> <td>2</td> </tr> </table>	Edible offal (mammalian)	Insert	0.03	Lettuce, head	Omit	T5		Substitute	5	Lettuce, leaf	Omit	T5		Substitute	7	Meat (mammalian) (in the fat)	Insert	0.05	Milk fats	Insert	0.05	Milks	Insert	*0.01	Peppers, Sweet	Omit	T1		Substitute	1	Tomato	Omit	T2		Substitute	2	<p>NEDI: 46% of the ADI</p>
Edible offal (mammalian)	Insert	0.03																																			
Lettuce, head	Omit	T5																																			
	Substitute	5																																			
Lettuce, leaf	Omit	T5																																			
	Substitute	7																																			
Meat (mammalian) (in the fat)	Insert	0.05																																			
Milk fats	Insert	0.05																																			
Milks	Insert	*0.01																																			
Peppers, Sweet	Omit	T1																																			
	Substitute	1																																			
Tomato	Omit	T2																																			
	Substitute	2																																			
<p>Flumetsulam Flumetsulam is a systemic herbicide. It is absorbed by roots and leaves of plants and translocated to growth points. It is used to control various weeds in cereals and pasture.</p> <p>The APVMA has approved a use pattern to control various weeds in pastures and cereal crops. No changes are proposed for cereal MRLs.</p> <table border="0" data-bbox="177 1727 983 1785"> <tr> <td>Edible offal (mammalian)</td> <td>Omit</td> <td>*0.2</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.3</td> </tr> </table>	Edible offal (mammalian)	Omit	*0.2		Substitute	0.3	<p>NEDI: <1% of the ADI</p>																														
Edible offal (mammalian)	Omit	*0.2																																			
	Substitute	0.3																																			

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																																																																																		
<p>Imazamox Imazamox is a imidazolinone herbicide. It is an acetolactate synthase (ALS) (also known as acetoxyacid synthase (AHAS)) inhibitor. It is absorbed through both foliage and roots and is translocated to growing points causing plants to wilt and turn brown. It is used for the early post emergent control of annual grass and broad leaf weeds in various crops.</p> <p>The APVMA has issued a permit for its use with imazapyr to control annual grass and broad leaf weeds in oilseed poppy. The recommended MRL is at the LOQ.</p> <p>Poppy seed Insert T*0.05</p>	<p>NEDI: <1% of the ADI</p>																																																																																																		
<p>Imazapyr Imazapyr is a systemic, contact and residual herbicide. It is absorbed by the foliage and roots and translocated via the xylem and phloem to the meristematic regions where it accumulates. It is used to control annual grass and broad leaf weeds in various crops.</p> <p>The APVMA has issued a permit for its use with imazamox to control annual grass and broad leaf weeds in oilseed poppy. The recommended MRL is at the LOQ.</p> <p>Poppy seed Insert T*0.05</p>	<p>NEDI: <1% of the ADI</p>																																																																																																		
<p>Imidacloprid Imidacloprid is a systemic insecticide with contact and stomach action. It acts on the central nervous system of insects causing blockage of postsynaptic nicotinic acetylcholine receptors. It is used as a seed dressing, or soil or foliar treatment to control sucking insects including aphids, thrips and whitefly in cereals, oilseeds, fruits and vegetables.</p> <p>The APVMA has approved its use as an in-furrow soil treatment to control green peach aphid and silverleaf whitefly on potatoes and sweet potatoes. The APVMA has evaluated further data in relation to the use of imidacloprid as a seedling or soil drench in certain leafy vegetables; and as a seed dressing to control aphids in broad beans, field peas and lentils. The data are sufficient to remove the temporary status of the recommended leafy vegetables MRLs. A lupin MRL is also recommended.</p> <table border="0" data-bbox="185 1541 975 2000"> <tr> <td>Broad bean (dry)</td> <td>Insert</td> <td>*0.05</td> <td></td> <td></td> </tr> <tr> <td>Field pea (dry)</td> <td>Insert</td> <td>*0.05</td> <td></td> <td></td> </tr> <tr> <td>Leafy vegetables [except lettuce, head]</td> <td>Insert</td> <td>20</td> <td></td> <td></td> </tr> <tr> <td>Leafy vegetables [except lettuce, leaf]</td> <td>Omit</td> <td>T5</td> <td></td> <td></td> </tr> <tr> <td>Lentil (dry)</td> <td>Insert</td> <td>0.2</td> <td><1</td> <td><1</td> </tr> <tr> <td>Lettuce, head</td> <td>Insert</td> <td>5</td> <td>12</td> <td>7</td> </tr> <tr> <td>Lettuce, leaf</td> <td>Omit</td> <td>T20</td> <td></td> <td></td> </tr> <tr> <td>Lupin (dry)</td> <td>Omit</td> <td>*0.05</td> <td></td> <td></td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.2</td> <td><1</td> <td><1</td> </tr> <tr> <td>Potato</td> <td>Omit</td> <td>T0.5</td> <td></td> <td></td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.3</td> <td><1</td> <td><1</td> </tr> <tr> <td>Sweet potato</td> <td>Omit</td> <td>0.5</td> <td></td> <td></td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.3</td> <td><1</td> <td><1</td> </tr> </table>	Broad bean (dry)	Insert	*0.05			Field pea (dry)	Insert	*0.05			Leafy vegetables [except lettuce, head]	Insert	20			Leafy vegetables [except lettuce, leaf]	Omit	T5			Lentil (dry)	Insert	0.2	<1	<1	Lettuce, head	Insert	5	12	7	Lettuce, leaf	Omit	T20			Lupin (dry)	Omit	*0.05				Substitute	0.2	<1	<1	Potato	Omit	T0.5				Substitute	0.3	<1	<1	Sweet potato	Omit	0.5				Substitute	0.3	<1	<1	<p>NEDI: 19% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="991 1518 1382 2000"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td>Broad bean (dry)</td> <td><1</td> <td><1</td> </tr> <tr> <td>Field pea (dry)</td> <td><1</td> <td><1</td> </tr> <tr> <td>Leafy vegetables [except lettuce, head]</td> <td>50</td> <td>36</td> </tr> <tr> <td>Lentil (dry)</td> <td><1</td> <td><1</td> </tr> <tr> <td>Lettuce, head</td> <td>12</td> <td>7</td> </tr> <tr> <td>Lettuce, leaf</td> <td></td> <td></td> </tr> <tr> <td>Lupin (dry)</td> <td></td> <td></td> </tr> <tr> <td>Potato</td> <td><1</td> <td><1</td> </tr> <tr> <td>Sweet potato</td> <td></td> <td></td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>	Broad bean (dry)	<1	<1	Field pea (dry)	<1	<1	Leafy vegetables [except lettuce, head]	50	36	Lentil (dry)	<1	<1	Lettuce, head	12	7	Lettuce, leaf			Lupin (dry)			Potato	<1	<1	Sweet potato				<1	<1
Broad bean (dry)	Insert	*0.05																																																																																																	
Field pea (dry)	Insert	*0.05																																																																																																	
Leafy vegetables [except lettuce, head]	Insert	20																																																																																																	
Leafy vegetables [except lettuce, leaf]	Omit	T5																																																																																																	
Lentil (dry)	Insert	0.2	<1	<1																																																																																															
Lettuce, head	Insert	5	12	7																																																																																															
Lettuce, leaf	Omit	T20																																																																																																	
Lupin (dry)	Omit	*0.05																																																																																																	
	Substitute	0.2	<1	<1																																																																																															
Potato	Omit	T0.5																																																																																																	
	Substitute	0.3	<1	<1																																																																																															
Sweet potato	Omit	0.5																																																																																																	
	Substitute	0.3	<1	<1																																																																																															
	<u>2-6 years</u>	<u>2+ years</u>																																																																																																	
Broad bean (dry)	<1	<1																																																																																																	
Field pea (dry)	<1	<1																																																																																																	
Leafy vegetables [except lettuce, head]	50	36																																																																																																	
Lentil (dry)	<1	<1																																																																																																	
Lettuce, head	12	7																																																																																																	
Lettuce, leaf																																																																																																			
Lupin (dry)																																																																																																			
Potato	<1	<1																																																																																																	
Sweet potato																																																																																																			
	<1	<1																																																																																																	

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment															
<p>Indoxacarb</p> <p>Indoxacarb is a foliar insecticide. It is active by contact and ingestion. It blocks sodium ion channels in nerve cells causing cessation of feeding, poor coordination, paralysis and death. It is used to control Lepidoptera in cotton, fruit and vegetables.</p> <p>The APVMA has issued a permit for its use to control Heliothis (<i>Helicoverpa armigera</i> and <i>Helicoverpa punctigera</i>) on peanuts.</p> <table> <tr> <td>Peanut</td> <td>Insert</td> <td>T0.02</td> </tr> </table>	Peanut	Insert	T0.02	<p>NEDI: 21% of the ADI</p> <p>NESTI as % of the ARfD</p> <table> <tr> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td><1</td> <td><1</td> </tr> </table>	<u>2-6 years</u>	<u>2+ years</u>	<1	<1								
Peanut	Insert	T0.02														
<u>2-6 years</u>	<u>2+ years</u>															
<1	<1															
<p>Iprodione</p> <p>Iprodione is a foliar fungicide with contact, protective and curative action. It inhibits spore germination and growth of fungal mycelium. It is used to control various moulds and rots including Sclerotinia (<i>Sclerotinia sclerotiorum</i>), grey mould (<i>Botrytis cinerea</i>) and Alternaria leaf spot (<i>Alternaria brassicae</i>) in cereals, oilseeds, pulses, nuts, fruits and vegetables.</p> <p>The APVMA has issued permits for its use to control sclerotinia rot in peppers and grey mould in Brussels sprouts. The data are sufficient to remove the temporary status of the Brussels sprouts MRL.</p> <table> <tr> <td>Brussels sprouts</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.5</td> </tr> <tr> <td>Peppers</td> <td>Insert</td> <td>T2</td> </tr> </table>	Brussels sprouts	Omit	T1		Substitute	0.5	Peppers	Insert	T2	<p>NEDI: 44% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: 1% of the ADI for adult males 25 – 34 years and toddlers 2 years and <1% of the ADI for other population groups assessed</p> <p>19th ATDS: 1% of the ADI for toddlers 2 years and <1% of the ADI for other population groups assessed</p>						
Brussels sprouts	Omit	T1														
	Substitute	0.5														
Peppers	Insert	T2														
<p>Lambda-cyhalothrin</p> <p>Lambda-cyhalothrin is a synthetic pyrethroid insecticide. It is a sodium channel modulator. It causes excessive stimulation of neurons by preventing the closure of voltage sensitive sodium channels. It is used to control a wide range of insect pests in cereal, fruit and vegetable crops.</p> <p>The NHC requested in its submission on MRL Proposal M1005 that FSANZ consider including an MRL for lambda-cyhalothrin residues in cherries in the Code harmonised with the United States MRL. Residues may occur in cherries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <p>Note: Lambda-cyhalothrin MRLs are listed under cyhalothrin.</p> <table> <tr> <td>Stone fruits</td> <td>Insert</td> <td>0.5</td> </tr> </table>	Stone fruits	Insert	0.5	<p>NEDI: 92% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results for cyhalothrin:</p> <p>20th ATDS: not detected in any foods sampled</p> <p>19th ATDS: <1% of the ADI for all population groups assessed</p> <p>NESTI as % of the ARfD</p> <table> <tr> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td>85</td> <td>Apricot 10</td> </tr> <tr> <td>41</td> <td>Cherries 7</td> </tr> <tr> <td>41</td> <td>Nectarine 23</td> </tr> <tr> <td>40</td> <td>Peach 16</td> </tr> <tr> <td>41</td> <td>Plum (including 15 prunes)</td> </tr> </table>	<u>2-6 years</u>	<u>2+ years</u>	85	Apricot 10	41	Cherries 7	41	Nectarine 23	40	Peach 16	41	Plum (including 15 prunes)
Stone fruits	Insert	0.5														
<u>2-6 years</u>	<u>2+ years</u>															
85	Apricot 10															
41	Cherries 7															
41	Nectarine 23															
40	Peach 16															
41	Plum (including 15 prunes)															

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																													
<p>Paclobutrazol Paclobutrazol is a plant growth regulator. It inhibits gibberellin and sterol synthesis. It is used on fruit trees to produce more compact plants (inhibit vegetative growth) and improve fruit set.</p> <p>The APVMA has issued a permit for its use on barley and wheat to reduce lodging.</p> <table border="0" data-bbox="177 504 983 562"> <tr> <td>Barley</td> <td>Insert</td> <td>T0.1</td> </tr> <tr> <td>Wheat</td> <td>Insert</td> <td>T0.1</td> </tr> </table>	Barley	Insert	T0.1	Wheat	Insert	T0.1	<p>NEDI: 15% of the ADI</p>																																							
Barley	Insert	T0.1																																												
Wheat	Insert	T0.1																																												
<p>Pendimethalin Pendimethalin is a selective herbicide. It is absorbed by the roots and leaves. It inhibits microtubule assembly. It is used to control annual grasses and broad leaf weeds in a wide range of crops.</p> <p>The APVMA has issued a permit for its use to control weeds in basil, bay trees, borage, chives, coriander, dill, fennel, lemon balm, lemon grass, kaffir lime, marigold, marjoram, oregano, mints, nasturtium, parsley, rosemary, sage, Burnet salad, sorrel, tarragon, savoury and thyme prior to transplanting. The data are sufficient to recommended an MRL at the LOQ.</p> <table border="0" data-bbox="177 929 983 965"> <tr> <td>Herbs</td> <td>Insert</td> <td>*0.05</td> </tr> </table>	Herbs	Insert	*0.05	<p>NEDI: <1% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: not detected in any foods sampled</p>																																										
Herbs	Insert	*0.05																																												
<p>Permethrin Permethrin is a non-systemic synthetic pyrethroid insecticide. It has contact and stomach action and a slight repellent effect. It acts on the nervous system of insects, disturbing the function of neurons by interaction with the sodium channel. It is used to control pests on a wide range of crops.</p> <p>The APVMA has issued a permit for its use to control Lepidopteran pests (including cabbage moth (<i>Helicoverpa</i> spp.), cluster caterpillar and cabbage white butterfly) on field grown basil, bay trees, borage, chives, coriander, dill, fennel, lemon balm, lemon grass, kaffir lime, marigold, marjoram, oregano, mints, nasturtium, parsley, rosemary, sage, Burnet salad, sorrel, tarragon, savoury and thyme.</p> <table border="0" data-bbox="177 1422 983 1706"> <tr> <td>Coriander (leaves and stems)</td> <td>Omit</td> <td>T10</td> </tr> <tr> <td>Coriander (leaves, stem, roots)</td> <td>Insert</td> <td>30</td> </tr> <tr> <td>Herbs</td> <td>Omit</td> <td>T10</td> </tr> <tr> <td></td> <td>Substitute</td> <td>30</td> </tr> <tr> <td>Kaffir lime leaves</td> <td>Omit</td> <td>T10</td> </tr> <tr> <td></td> <td>Substitute</td> <td>30</td> </tr> <tr> <td>Lemon balm</td> <td>Insert</td> <td>30</td> </tr> <tr> <td>Lemon grass</td> <td>Omit</td> <td>T10</td> </tr> <tr> <td></td> <td>Substitute</td> <td>30</td> </tr> </table>	Coriander (leaves and stems)	Omit	T10	Coriander (leaves, stem, roots)	Insert	30	Herbs	Omit	T10		Substitute	30	Kaffir lime leaves	Omit	T10		Substitute	30	Lemon balm	Insert	30	Lemon grass	Omit	T10		Substitute	30	<p>NEDI: 17% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>19th ATDS: <1% of the ADI for all population groups assessed</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1388 1390 1706"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> <tr> <td></td> <td><1</td> <td><1</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>		<1	<1		<1	<1		<1	<1		<1	<1		<1	<1
Coriander (leaves and stems)	Omit	T10																																												
Coriander (leaves, stem, roots)	Insert	30																																												
Herbs	Omit	T10																																												
	Substitute	30																																												
Kaffir lime leaves	Omit	T10																																												
	Substitute	30																																												
Lemon balm	Insert	30																																												
Lemon grass	Omit	T10																																												
	Substitute	30																																												
	<u>2-6 years</u>	<u>2+ years</u>																																												
	<1	<1																																												
	<1	<1																																												
	<1	<1																																												
	<1	<1																																												
	<1	<1																																												

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																																
<p>Phosphorous acid Phosphorous acid is a selective systemic phosphonate fungicide with multi-site activity. It creates an immune response within the host plant and also has direct antifungal activity. It is used to control fungal diseases on fruits and vegetables.</p> <p>The APVMA has issued permits for its use to control <i>Pythium</i> rhizome rot (<i>Pythium myriotylum</i>) in ginger and <i>Phytophthora</i> root rot (<i>Phytophthora nicotianae</i> var. <i>nicotianae</i>) in field grown tomatoes grown for processing purposes.</p> <table border="0" data-bbox="177 593 983 656"> <tr> <td>Ginger, root</td> <td>Insert</td> <td>T100</td> </tr> <tr> <td>Tomato</td> <td>Insert</td> <td>T100</td> </tr> </table>	Ginger, root	Insert	T100	Tomato	Insert	T100	<p>NEDI: 7% of the ADI</p>																																										
Ginger, root	Insert	T100																																															
Tomato	Insert	T100																																															
<p>Pirimicarb Pirimicarb is a selective systemic insecticide. It has contact, stomach and respiratory action. It is an anticholinesterase inhibitor. It is used to control aphids on crops and pastures.</p> <p>The APVMA has issued permits for its use to control aphid, including green aphids and cabbage aphids on leafy vegetables; aphids on spring onions; and cowpea aphid (<i>Aphis craccivora</i>) and soya bean aphid (<i>Aphis glycines</i>) on adzuki bean, mung bean and soy bean. MRLs are also recommended for shallots and Welsh onions. Shallots may be harvested green and referred to as spring onions and Welsh onions may be considered to be spring onions. The current soy bean MRL remains appropriate.</p> <table border="0" data-bbox="177 1288 983 1872"> <tr> <td>Adzuki bean (dry)</td> <td>Insert</td> <td>T0.5</td> </tr> <tr> <td>Leafy vegetables [except chervil; mizuna; rucola]</td> <td>Omit</td> <td>T5</td> </tr> <tr> <td>Leafy vegetables [except chervil; mizuna; rucola (rocket)]</td> <td>Insert</td> <td>T7</td> </tr> <tr> <td>Mung bean (dry)</td> <td>Insert</td> <td>T0.5</td> </tr> <tr> <td>Onion, Welsh</td> <td>Insert</td> <td>T3</td> </tr> <tr> <td>Shallot</td> <td>Insert</td> <td>T3</td> </tr> <tr> <td>Spring onion</td> <td>Insert</td> <td>T3</td> </tr> <tr> <td>Vegetables [except leafy vegetables; lupin (dry); soya bean (dry); sweet corn (corn-on-the-cob)]</td> <td>Omit</td> <td>1</td> </tr> <tr> <td>Vegetables [except adzuki bean (dry); leafy vegetables; lupin (dry); mung bean (dry); onion, Welsh; shallot; soya bean (dry); spring onion; sweet corn (corn-on-the-cob)]</td> <td>Insert</td> <td>1</td> </tr> </table>	Adzuki bean (dry)	Insert	T0.5	Leafy vegetables [except chervil; mizuna; rucola]	Omit	T5	Leafy vegetables [except chervil; mizuna; rucola (rocket)]	Insert	T7	Mung bean (dry)	Insert	T0.5	Onion, Welsh	Insert	T3	Shallot	Insert	T3	Spring onion	Insert	T3	Vegetables [except leafy vegetables; lupin (dry); soya bean (dry); sweet corn (corn-on-the-cob)]	Omit	1	Vegetables [except adzuki bean (dry); leafy vegetables; lupin (dry); mung bean (dry); onion, Welsh; shallot; soya bean (dry); spring onion; sweet corn (corn-on-the-cob)]	Insert	1	<p>NEDI: 89% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>19th ATDS: <1% of the ADI for all population groups assessed</p> <p>Note that the proposed vegetables MRL variation is a technical amendment only. NESTI calculations are not required.</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1265 1390 1344"> <thead> <tr> <th></th> <th><u>2-6 years</u></th> <th><u>2+ years</u></th> </tr> </thead> <tbody> <tr> <td></td> <td>10</td> <td>2</td> </tr> </tbody> </table> <table border="0" data-bbox="983 1377 1390 1579"> <tbody> <tr> <td></td> <td>64</td> <td>33</td> </tr> <tr> <td></td> <td>10</td> <td>2</td> </tr> <tr> <td></td> <td>15</td> <td>3</td> </tr> <tr> <td></td> <td>11</td> <td>2</td> </tr> <tr> <td></td> <td>7</td> <td>2</td> </tr> </tbody> </table>		<u>2-6 years</u>	<u>2+ years</u>		10	2		64	33		10	2		15	3		11	2		7	2
Adzuki bean (dry)	Insert	T0.5																																															
Leafy vegetables [except chervil; mizuna; rucola]	Omit	T5																																															
Leafy vegetables [except chervil; mizuna; rucola (rocket)]	Insert	T7																																															
Mung bean (dry)	Insert	T0.5																																															
Onion, Welsh	Insert	T3																																															
Shallot	Insert	T3																																															
Spring onion	Insert	T3																																															
Vegetables [except leafy vegetables; lupin (dry); soya bean (dry); sweet corn (corn-on-the-cob)]	Omit	1																																															
Vegetables [except adzuki bean (dry); leafy vegetables; lupin (dry); mung bean (dry); onion, Welsh; shallot; soya bean (dry); spring onion; sweet corn (corn-on-the-cob)]	Insert	1																																															
	<u>2-6 years</u>	<u>2+ years</u>																																															
	10	2																																															
	64	33																																															
	10	2																																															
	15	3																																															
	11	2																																															
	7	2																																															

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																													
<p>Pyraclostrobin Pyraclostrobin is a strobiluran fungicide with protectant, curative and translaminar properties. It inhibits mitochondrial respiration by blocking electron transfer within the respiratory chain; this severely disrupts cellular biochemical processes and results in cessation of fungal growth. It is used to control major plant pathogens in fruit and vegetables.</p> <p>The APVMA has issued permits for its use to control black spot (<i>Asperisporium caricae</i>) and brown spot (<i>Corynespora cassiicola</i>) on pawpaw and Pseudocercospora leaf spot (<i>P. Anonicola</i>) on custard apple. The APVMA has also approved its use with epoxiconazole to control various diseases in wheat, barley and oats. The recommended cereal grains MRL is at the LOQ.</p> <table border="0" data-bbox="177 741 983 869"> <tr> <td>Cereal grains</td> <td>Insert</td> <td>*0.01</td> </tr> <tr> <td>Custard apple</td> <td>Insert</td> <td>T3</td> </tr> <tr> <td>Papaya (pawpaw)</td> <td>Insert</td> <td>T0.5</td> </tr> </table>	Cereal grains	Insert	*0.01	Custard apple	Insert	T3	Papaya (pawpaw)	Insert	T0.5	<p>NEDI: 3% of the ADI</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 712 1390 869"> <tr> <td></td> <td><u>2-6 years</u></td> <td></td> <td><u>2+ years</u></td> </tr> <tr> <td></td> <td><1</td> <td></td> <td><1</td> </tr> <tr> <td></td> <td>48</td> <td>Tropical fruit</td> <td>13</td> </tr> <tr> <td></td> <td></td> <td>inedible peel</td> <td></td> </tr> <tr> <td></td> <td>11</td> <td>Pineapple</td> <td>4</td> </tr> </table>		<u>2-6 years</u>		<u>2+ years</u>		<1		<1		48	Tropical fruit	13			inedible peel			11	Pineapple	4
Cereal grains	Insert	*0.01																												
Custard apple	Insert	T3																												
Papaya (pawpaw)	Insert	T0.5																												
	<u>2-6 years</u>		<u>2+ years</u>																											
	<1		<1																											
	48	Tropical fruit	13																											
		inedible peel																												
	11	Pineapple	4																											
<p>Pyrimethanil Pyrimethanil is a foliar fungicide with protectant action. It inhibits fungal enzymes necessary for infection. It is used to control fungal diseases in a range of horticultural situations.</p> <p>The APVMA has issued a permit for its use with chlorothalonil to control Alternaria and Botrytis on chickory, endive, radicchio, silverbeet and spinach.</p> <table border="0" data-bbox="177 1234 983 1263"> <tr> <td>Leafy vegetables</td> <td>Insert</td> <td>T5</td> </tr> </table>	Leafy vegetables	Insert	T5	<p>NEDI: 5% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: <1% of the ADI for all population groups assessed</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="983 1205 1390 1263"> <tr> <td></td> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td></td> <td>5</td> <td>3</td> </tr> </table>		<u>2-6 years</u>	<u>2+ years</u>		5	3																				
Leafy vegetables	Insert	T5																												
	<u>2-6 years</u>	<u>2+ years</u>																												
	5	3																												
<p>Pyriproxifen Pyriproxifen is an insecticide. It is an insect growth regulator, which inhibits metamorphosis and reproduction. It is used to control silverleaf whitefly in cotton; silverleaf whitefly and greenhouse whitefly in cucurbits, tomatoes and eggplant; and various scale insects in citrus fruit, mangoes, olives, coffee and passionfruit.</p> <p>The APVMA has approved a use pattern to control pests in mango.</p> <table border="0" data-bbox="177 1603 983 1659"> <tr> <td>Mango</td> <td>Omit</td> <td>*0.01</td> </tr> <tr> <td></td> <td>Substitute</td> <td>0.05</td> </tr> </table>	Mango	Omit	*0.01		Substitute	0.05	<p>NEDI: 2% of the ADI</p>																							
Mango	Omit	*0.01																												
	Substitute	0.05																												

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																		
<p>Simazine Simazine is a selective systemic herbicide. It is absorbed principally through the roots but also through foliage, with translocation acropetally in the xylem accumulating in the apical meristems and leaves. It inhibits photosynthetic electron transport.</p> <p>The APVMA has issued a permit for its use to control blue green algae in dams, tanks and troughs for livestock watering. The recommended MRLs are at the LOQ.</p> <table border="0" data-bbox="177 593 986 770"> <tr> <td data-bbox="177 593 606 627">Edible offal (mammalian)</td> <td data-bbox="606 593 893 627">Omit</td> <td data-bbox="893 593 986 627">*0.01</td> </tr> <tr> <td></td> <td data-bbox="606 627 893 660">Substitute</td> <td data-bbox="893 627 986 660">*0.05</td> </tr> <tr> <td data-bbox="177 660 606 694">Meat (mammalian)</td> <td data-bbox="606 660 893 694">Omit</td> <td data-bbox="893 660 986 694">*0.01</td> </tr> <tr> <td></td> <td data-bbox="606 694 893 728">Substitute</td> <td data-bbox="893 694 986 728">*0.05</td> </tr> <tr> <td data-bbox="177 728 606 761">Milks</td> <td data-bbox="606 728 893 761">Omit</td> <td data-bbox="893 728 986 761">*0.01</td> </tr> <tr> <td></td> <td data-bbox="606 761 893 770">Substitute</td> <td data-bbox="893 761 986 770">*0.02</td> </tr> </table>	Edible offal (mammalian)	Omit	*0.01		Substitute	*0.05	Meat (mammalian)	Omit	*0.01		Substitute	*0.05	Milks	Omit	*0.01		Substitute	*0.02	<p>NEDI: 16% of the ADI</p>
Edible offal (mammalian)	Omit	*0.01																	
	Substitute	*0.05																	
Meat (mammalian)	Omit	*0.01																	
	Substitute	*0.05																	
Milks	Omit	*0.01																	
	Substitute	*0.02																	

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment		
<p>Spirotetramat</p>	NEDI: 12% of the ADI		
<p>Spirotetramat is a cyclic ketoenole insecticide. It is a tetramic acid derivative. It inhibits acetyl CoA carboxylase, a key enzyme in fatty acid biosynthesis. It is active against a wide spectrum of sucking insects including aphids, scales, mealybugs, whiteflies, psyllids and certain thrips.</p>			
<p>The APVMA has approved an extension of its use to control various pests in brassicas, cucurbits, eggplant, capsicum, chillies, tomatoes, potatoes, sweet potatoes and leafy vegetables. The APVMA has evaluated further trial data in relation to use of spirotetramat to control pests in citrus fruits, mango and onion. The data are sufficient to remove the temporary status of the MRLs. The APVMA has also issued a permit for use of spirotetramat to control various pests on beans and peas.</p>			
<p>Bayer requested MRLs in the Code harmonised with the Codex MRLs for spirotetramat residues in grapes and raisins. Residues may occur in imported grapes and raisins. The MRL may minimise potential trade disruption and extend consumer choice.</p>			
	NESTI as % of the ARfD		
	<u>2-6 years</u>	<u>2+ years</u>	
Citrus fruits	Omit	T1	
	Substitute	1	5 2
Dried grapes	Insert	4	<1 <1
Fruiting vegetables, cucurbits	Omit	T2	
Fruiting vegetables, cucurbits	Insert	2	2 2 Zucchini <1
[except melons]			<1 Cucumber <1
Fruiting vegetables, other than cucurbits	Insert	7	12 7
Grapes	Insert	2	14 3
Leafy vegetables [except lettuce, head]	Insert	5	3 2 Lettuce, leaf 2
Legume vegetables	Insert	T2	<1 2 Spinach 2
Lettuce, head	Omit	T5	<1 <1
	Substitute	3	<1 <1
Lettuce, leaf	Omit	T10	
Mango	Omit	T0.3	
	Substitute	0.3	1 <1
Melons, except watermelon	Insert	0.5	1 <1
Onion, bulb	Omit	T0.5	
	Substitute	0.5	<1 <1
Peppers, Sweet	Omit	T5	
Potato	Insert	5	6 2
Sweet potato	Insert	5	2 2
Tomato	Omit	T7	
Watermelon	Insert	0.5	1 1

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment				
<p>Tebuconazole Tebuconazole is a non-systemic foliar triazole fungicide. It has protective action. It inhibits steroid demethylation leading to inhibition of ergosterol biosynthesis. It is used to control various fungal diseases in many crops.</p> <p>The NHC requested in its submission on MRL Proposal M1005 that FSANZ consider including an MRL for tebuconazole residues in cherries in the Code harmonised with the United States MRL. Residues may occur in cherries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <p>Cherries Insert 5</p>	<p>NEDI: 24% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: not detected in any foods sampled</p>				
<p>Tebufenozide Tebufenozide is an ecdysone agonist insecticide. It binds to the receptor site of the insect moulting hormone ecdysone. It lethally accelerates the moulting process. It is used to control Lepidopteran larvae on fruits nuts and other crops.</p> <p>The CMC requested that FSANZ include an MRL in the Code harmonised with the Codex limit for tebufenozide residues in cranberries. Residues may occur in cranberries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <p>Cranberry Insert 0.5</p>	<p>NEDI: 32% of the ADI</p> <p>NESTI as % of the ARfD</p> <table style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;"><u>2-6 years</u></td> <td style="text-align: center;"><u>2+ years</u></td> </tr> <tr> <td style="text-align: center;"><1</td> <td style="text-align: center;"><1</td> </tr> </table>	<u>2-6 years</u>	<u>2+ years</u>	<1	<1
<u>2-6 years</u>	<u>2+ years</u>				
<1	<1				

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																																																									
<p>Terbuthylazine Terbuthylazine is a herbicide. It is absorbed mainly by the roots. It inhibits photosynthetic electron transport at the photosystem II receptor site.</p> <p>The APVMA has evaluated further data in relation to the approved use of spirotetramat to control a wide variety of weeds in pre-emergent lupins, chickpeas, field peas, fava beans and certain canola varieties. The data are sufficient to confirm the MRLs. The APVMA has also issued a permit for use of spirotetramat to control various weeds in sorghum, maize and sweet corn. The recommended MRLs are at the LOQ.</p> <table border="0"> <tr> <td data-bbox="177 656 619 685">Edible offal (mammalian)</td> <td data-bbox="619 656 879 685">Omit</td> <td data-bbox="879 656 983 685">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 685 879 714">Substitute</td> <td data-bbox="879 685 983 714">*0.01</td> </tr> <tr> <td data-bbox="177 714 619 743">Eggs</td> <td data-bbox="619 714 879 743">Omit</td> <td data-bbox="879 714 983 743">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 743 879 772">Substitute</td> <td data-bbox="879 743 983 772">*0.01</td> </tr> <tr> <td data-bbox="177 772 619 801">Maize</td> <td data-bbox="619 772 879 801">Insert</td> <td data-bbox="879 772 983 801">T*0.02</td> </tr> <tr> <td data-bbox="177 801 619 831">Meat (mammalian)</td> <td data-bbox="619 801 879 831">Omit</td> <td data-bbox="879 801 983 831">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 831 879 860">Substitute</td> <td data-bbox="879 831 983 860">*0.01</td> </tr> <tr> <td data-bbox="177 860 619 889">Milks</td> <td data-bbox="619 860 879 889">Omit</td> <td data-bbox="879 860 983 889">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 889 879 918">Substitute</td> <td data-bbox="879 889 983 918">*0.01</td> </tr> <tr> <td data-bbox="177 918 619 947">Poultry, edible offal of</td> <td data-bbox="619 918 879 947">Omit</td> <td data-bbox="879 918 983 947">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 947 879 976">Substitute</td> <td data-bbox="879 947 983 976">*0.01</td> </tr> <tr> <td data-bbox="177 976 619 1005">Poultry meat</td> <td data-bbox="619 976 879 1005">Omit</td> <td data-bbox="879 976 983 1005">T*0.01</td> </tr> <tr> <td></td> <td data-bbox="619 1005 879 1034">Substitute</td> <td data-bbox="879 1005 983 1034">*0.01</td> </tr> <tr> <td data-bbox="177 1034 619 1064">Pulses</td> <td data-bbox="619 1034 879 1064">Omit</td> <td data-bbox="879 1034 983 1064">T*0.02</td> </tr> <tr> <td></td> <td data-bbox="619 1064 879 1093">Substitute</td> <td data-bbox="879 1064 983 1093">*0.02</td> </tr> <tr> <td data-bbox="177 1093 619 1122">Rape seed (canola)</td> <td data-bbox="619 1093 879 1122">Omit</td> <td data-bbox="879 1093 983 1122">T*0.02</td> </tr> <tr> <td></td> <td data-bbox="619 1122 879 1151">Substitute</td> <td data-bbox="879 1122 983 1151">*0.02</td> </tr> <tr> <td data-bbox="177 1151 619 1180">Sorghum</td> <td data-bbox="619 1151 879 1180">Insert</td> <td data-bbox="879 1151 983 1180">T*0.02</td> </tr> <tr> <td data-bbox="177 1180 619 1209">Sweet corn (corn-on-the-cob)</td> <td data-bbox="619 1180 879 1209">Insert</td> <td data-bbox="879 1180 983 1209">T*0.02</td> </tr> </table>	Edible offal (mammalian)	Omit	T*0.01		Substitute	*0.01	Eggs	Omit	T*0.01		Substitute	*0.01	Maize	Insert	T*0.02	Meat (mammalian)	Omit	T*0.01		Substitute	*0.01	Milks	Omit	T*0.01		Substitute	*0.01	Poultry, edible offal of	Omit	T*0.01		Substitute	*0.01	Poultry meat	Omit	T*0.01		Substitute	*0.01	Pulses	Omit	T*0.02		Substitute	*0.02	Rape seed (canola)	Omit	T*0.02		Substitute	*0.02	Sorghum	Insert	T*0.02	Sweet corn (corn-on-the-cob)	Insert	T*0.02	<p>NEDI: 4% of the ADI</p>
Edible offal (mammalian)	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Eggs	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Maize	Insert	T*0.02																																																								
Meat (mammalian)	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Milks	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Poultry, edible offal of	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Poultry meat	Omit	T*0.01																																																								
	Substitute	*0.01																																																								
Pulses	Omit	T*0.02																																																								
	Substitute	*0.02																																																								
Rape seed (canola)	Omit	T*0.02																																																								
	Substitute	*0.02																																																								
Sorghum	Insert	T*0.02																																																								
Sweet corn (corn-on-the-cob)	Insert	T*0.02																																																								
<p>Tolclofos-methyl Tolclofos-methyl is a non-systemic nitrophenyl fungicide with contact, protective and curative action. It is used as a seed or in-furrow treatment to control fungal diseases in beetroot, cotton and potatoes.</p> <p>The APVMA has issued a permit for its use to control <i>Rhizoctonia</i> fungi in beetroot and potato. The established potato MRL remains appropriate. The recommended MRL is at the LOQ.</p> <table border="0"> <tr> <td data-bbox="177 1543 619 1572">Beetroot</td> <td data-bbox="619 1543 879 1572">Omit</td> <td data-bbox="879 1543 983 1572">T0.5</td> </tr> <tr> <td></td> <td data-bbox="619 1572 879 1601">Substitute</td> <td data-bbox="879 1572 983 1601">*0.01</td> </tr> </table>	Beetroot	Omit	T0.5		Substitute	*0.01	<p>NEDI: <1% of the ADI</p>																																																			
Beetroot	Omit	T0.5																																																								
	Substitute	*0.01																																																								

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment															
<p>Triadimenol Triadimenol is a systemic fungicide with protective, curative and eradicator action. It is absorbed by roots and leaves, with ready translocation in young growing tissues, but less ready translocation in older, woody tissues. It inhibits gibberellin and ergosterol biosynthesis and hence the rate of cell division. It is used to control various fungal diseases in a range of crops.</p> <p>The APVMA has issued a permit for its use to control fungal diseases on tomatoes, eggplant, capsicums and chillies. Established MRLs for capsicum, tomato and eggplant remain appropriate. A temporary MRL is recommended for peppers, this group includes capsicum (sweet peppers) and chillies.</p> <table border="0" data-bbox="188 712 970 779"> <tr> <td>Peppers, Sweet</td> <td>Omit</td> <td>T1</td> </tr> <tr> <td>Peppers</td> <td>Insert</td> <td>T1</td> </tr> </table>	Peppers, Sweet	Omit	T1	Peppers	Insert	T1	<p>NEDI: 2% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: not detected in any foods sampled</p> <p>19th ATDS: not detected in any foods sampled</p> <p>NESTI as % of the ARfD</p> <table border="0" data-bbox="994 683 1348 806"> <tr> <td></td> <td><u>2-6 years</u></td> <td><u>2+ years</u></td> </tr> <tr> <td>12</td> <td>Capsicum</td> <td>5</td> </tr> <tr> <td>10[†]</td> <td>Chillies</td> <td>1</td> </tr> </table> <p>† Calculated using consumption data for capsicum as there is insufficient chilli consumption data for this age group.</p>		<u>2-6 years</u>	<u>2+ years</u>	12	Capsicum	5	10 [†]	Chillies	1
Peppers, Sweet	Omit	T1														
Peppers	Insert	T1														
	<u>2-6 years</u>	<u>2+ years</u>														
12	Capsicum	5														
10 [†]	Chillies	1														
<p>Trichlorfon Trichlorfon is an organophosphate insecticide. It binds irreversibly to the active site of acetylcholinesterase. Acetylcholinesterase is inactivated and therefore normal nerve impulse transmission is affected and the insect is paralysed. Trichlorfon is used to control pests in agriculture, horticulture, aquaculture and livestock.</p> <p>The APVMA has issued a permit for its use to control and treat skin and gill flukes, anchor worm (<i>Lernaea</i> spp.) and gill maggots (<i>Ergasilus</i> spp.) in farmed silver perch. Detectable residues are not expected to occur. The recommended MRL is at the LOQ.</p> <p>The proposed fruit and vegetables MRL variations are technical amendments.</p> <table border="0" data-bbox="188 1422 970 1785"> <tr> <td>Fish muscle</td> <td>Insert</td> <td>T*0.01</td> </tr> <tr> <td>Fruit [except as otherwise listed under this chemical]</td> <td>Omit</td> <td>0.1</td> </tr> <tr> <td>Fruit [except banana; dried fruits; peach]</td> <td>Insert</td> <td>0.1</td> </tr> <tr> <td>Vegetables [except as otherwise listed under this chemical]</td> <td>Omit</td> <td>0.1</td> </tr> <tr> <td>Vegetables [except beetroot; Brussels sprouts; cauliflower; celery; kale; peppers; pulses; sugar beet; sweet corn (corn-on-the-cob)]</td> <td>Insert</td> <td>0.1</td> </tr> </table>	Fish muscle	Insert	T*0.01	Fruit [except as otherwise listed under this chemical]	Omit	0.1	Fruit [except banana; dried fruits; peach]	Insert	0.1	Vegetables [except as otherwise listed under this chemical]	Omit	0.1	Vegetables [except beetroot; Brussels sprouts; cauliflower; celery; kale; peppers; pulses; sugar beet; sweet corn (corn-on-the-cob)]	Insert	0.1	<p>NEDI: 99% of the ADI</p> <p>Mean estimated daily dietary exposure based on mean analytical results:</p> <p>20th ATDS: not detected in any foods sampled</p>
Fish muscle	Insert	T*0.01														
Fruit [except as otherwise listed under this chemical]	Omit	0.1														
Fruit [except banana; dried fruits; peach]	Insert	0.1														
Vegetables [except as otherwise listed under this chemical]	Omit	0.1														
Vegetables [except beetroot; Brussels sprouts; cauliflower; celery; kale; peppers; pulses; sugar beet; sweet corn (corn-on-the-cob)]	Insert	0.1														

Requested MRLs expressed in milligrams of the chemical per kilogram of the food (mg/kg)	Dietary Exposure Assessment																		
<p>Trifloxystrobin Trifloxystrobin is a mesostematic, broad-spectrum fungicide with preventative and specific curative action. It inhibits mitochondrial respiration by blocking electron transfer at the Qo centre of cytochrome bc1. It is used to control powdery mildew, leaf spot and rust in horticultural situations.</p> <p>The APVMA has issued permits for its use to control <i>Cercospora</i> leaf spot (<i>Cercospora apii</i>) and <i>Septoria</i> spot (<i>Septoria apiicola</i>) in celery and powdery mildew in field grown silver beet, chicory, spinach and endive.</p> <p>The NHC requested that FSANZ consider including an MRL for trifloxystrobin residues in cherries in the Code harmonised with the United States MRL. Residues may occur in cherries imported from the United States. The MRL may minimise potential trade disruption and extend consumer choice.</p> <table border="0" data-bbox="188 801 971 987"> <tr> <td>Celery</td> <td>Insert</td> <td>T1</td> </tr> <tr> <td>Chard (silver beet)</td> <td>Insert</td> <td>T0.7</td> </tr> <tr> <td>Chicory leaves</td> <td>Insert</td> <td>T0.7</td> </tr> <tr> <td>Endive</td> <td>Insert</td> <td>T0.7</td> </tr> <tr> <td>Spinach</td> <td>Insert</td> <td>T0.7</td> </tr> <tr> <td>Stone fruits</td> <td>Insert</td> <td>2</td> </tr> </table>	Celery	Insert	T1	Chard (silver beet)	Insert	T0.7	Chicory leaves	Insert	T0.7	Endive	Insert	T0.7	Spinach	Insert	T0.7	Stone fruits	Insert	2	<p>NEDI: 4% of the ADI</p>
Celery	Insert	T1																	
Chard (silver beet)	Insert	T0.7																	
Chicory leaves	Insert	T0.7																	
Endive	Insert	T0.7																	
Spinach	Insert	T0.7																	
Stone fruits	Insert	2																	
<p>Trifluralin Trifluralin is a selective soil herbicide. It disrupts cell division and root development. It is applied to the soil and enters the seedling in the hypocotyl region. It is used for the pre-emergent control of broad leaf and annual grass weeds in a wide range of crops and horticultural situations.</p> <p>The APVMA has issued a permit for its use, pre-planting, to control certain weeds in chia. The recommended MRL is at the LOQ.</p> <table border="0" data-bbox="188 1328 971 1357"> <tr> <td>Chia</td> <td>Insert</td> <td>T*0.01</td> </tr> </table>	Chia	Insert	T*0.01	<p>NEDI: 7% of the ADI</p>															
Chia	Insert	T*0.01																	
<p>Trinexapac-ethyl Trinexapac-ethyl is a plant growth regulator and retardant. It is an internode elongation disruptor. It is absorbed by the foliage and translocated to the growing shoot. It is used to increase seed set, alkaloid levels and yield; and prevent lodging and stem elongation in sugar cane.</p> <p>The APVMA has issued a permit for its use in barley and wheat to reduce lodging.</p> <table border="0" data-bbox="188 1666 971 1727"> <tr> <td>Barley</td> <td>Insert</td> <td>T0.3</td> </tr> <tr> <td>Wheat</td> <td>Insert</td> <td>T0.3</td> </tr> </table>	Barley	Insert	T0.3	Wheat	Insert	T0.3	<p>NEDI: 3% of the ADI</p>												
Barley	Insert	T0.3																	
Wheat	Insert	T0.3																	